

**ERROR ESTIMATES FOR A CLASS OF FINITE  
DIFFERENCE-QUADRATURE SCHEMES FOR FULLY  
NONLINEAR DEGENERATE PARABOLIC INTEGRO-PDES**

I. H. BISWAS, E. R. JAKOBSEN, AND K. H. KARLSEN

ABSTRACT. Error estimates are derived for a class of monotone finite difference-quadrature schemes approximating viscosity solutions of nonlinear degenerate parabolic integro-PDEs with variable diffusion coefficients. The relevant equations can be viewed as Bellman equations associated to a class of controlled jump-diffusion (Lévy) processes. Our results cover both finite and infinite activity cases.

1. INTRODUCTION

In this article we consider error estimates for finite difference type numerical schemes for degenerate and fully nonlinear parabolic integro-partial differential equations (integro-PDEs henceforth) of Bellman type. We write the equation in the following abstract form,

$$(1.1) \quad u_t(t, x) + F(t, x, u(t, x), Du(t, x), D^2u(t, x), u(t, \cdot)) = 0 \text{ in } Q_T$$

where  $T > 0$  is a constant and  $Q_T = [0, T) \times \mathbb{R}^d$  and we impose a terminal condition,

$$(1.2) \quad u(T, x) = u_0(x) \text{ for all } x \in \mathbb{R}^d.$$

The nonlocal feature of the equation is indicated by the term  $u(t, \cdot)$ . For any  $(t, x, r, p, X) \in \mathbb{R} \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d$  and for any ‘sufficiently well behaved’  $\varphi$ , the nonlinearity  $F$  is defined as follows

$$F(t, x, r, p, X, \varphi(\cdot)) = \sup_{\alpha \in \mathcal{A}} \left\{ \frac{1}{2} \text{tr} [a^\alpha(t, x)X] + b^\alpha(t, x) \cdot p + \mathcal{I}^\alpha \varphi - c^\alpha(t, x)r + f^\alpha(t, x) \right\},$$

where the integral operator  $\mathcal{I}^\alpha$  is defined as

$$(1.3) \quad \begin{aligned} \mathcal{I}^\alpha \varphi(t, x) &= \int_E [\varphi(t, x + \eta^\alpha(x, z)) - \varphi(t, x) - \mathbf{1}_{|z| < 1} \eta^\alpha(x, z) \cdot D\varphi(t, x)] \nu(dz), \end{aligned}$$

---

*Date:* March 5, 2008.

*2000 Mathematics Subject Classification.* 45K05, 49L25, 65M12, 65L70.

*Key words and phrases.* Integro-partial differential equation, viscosity solution, finite difference scheme, error estimate, stochastic optimal control, Lévy process, Bellman equation.

This research is supported by the Research Council of Norway through an Outstanding Young Investigators Award (KHK) and partially through the project ‘Integro-PDEs: Numerical methods, Analysis, and Applications to Finance’.

and  $E = \mathbb{R}^M \setminus \{0\}$  ( $M$  integer) and  $\nu(dz)$  is a positive Radon measure on  $E$  – the so-called Lévy measure possessing at most a second order singularity at the origin and typically exponential decay at infinity.

The set  $\mathcal{A}$ , the value set of all admissible controls, is a compact metric space and the coefficients  $a^\alpha, \eta^\alpha, b^\alpha, c^\alpha, f^\alpha, u_0$  are sufficiently regular functions taking values in  $\mathbb{R}^{d \times d}, \mathbb{R}^d, \mathbb{R}^d, \mathbb{R}, \mathbb{R}, \mathbb{R}$  respectively. In this paper we will need  $F$  to have special structure. The precise structure and assumptions on the coefficients will be given in the next section.

Equation (1.1) is *degenerate parabolic* since we will allow (i) the diffusion matrices  $a^\alpha(t, x)$  merely to be non-negative definite and (ii) the jump vector  $\eta^\alpha(x, z)$  to be zero for some  $\alpha, x, z$ . In other words there is no (global) regularization in this problem, neither from the second derivative (“Laplacian smoothing”) nor from the integral term (“fractional Laplacian smoothing”). In general equation (1.1) will therefore not have classical solutions. For the type nonlinearity and degeneracy present in (1.1) it is natural to interpret solutions in the viscosity sense. The viscosity solution theory for the second order nonlinear partial differential equations is now well developed and has become an essential tool to study the optimal control problems for pure diffusion processes. In the past few years, there has been a considerable effort to extend the theory of viscosity solution to the integro-PDEs [1, 2, 3, 8, 9, 10, 16, 17, 22]. Although this theory is not as developed as its pure PDE counter part, it is good enough to provide existence, uniqueness, comparison principles, and some regularity results in certain situations.

Although this connection will not be exploited herein, equations of the form (1.1) appear as the Bellman equations associated to the optimal control of jump-diffusion processes (or Lévy processes) over a finite time horizon (see [22, 23]). Examples include various types of portfolio optimization problems in which the risky asset is driven by a Lévy process. The linear version of (1.1) is of particular relevance to pricing theory of European option. For more information on pricing theory and its relation to linear integro-partial differential equations, we refer to [12].

In this paper we focus on finite difference-quadrature type schemes for (1.1) and their convergence properties. To be more precise, we will derive error estimates for numerical schemes for non-local equations of the form (1.1). There is a considerable literature addressing the issue of convergence of approximate (numerical) solutions to second order PDEs in the viscosity solution framework, see for example [7, 13, 14]. The question of error estimate for numerical schemes, including finite difference type, is much more difficult and remained open until the recent works by Krylov [21, 19, 20] and Barles & Jakobsen [4, 5, 6, 15].

On the other hand, finding error estimate for approximation schemes for fully nonlinear integro-PDEs is largely an untouched area with very few published results. In a recent development [18]; Jakobsen, Karlsen and La Chioma have given a general framework for proving error estimates in the stationary case. To apply this framework strong assumptions are placed on the schemes, and in [18] they are verified only when the diffusion matrices  $a^\alpha$  is independent of the space variable  $x$ . In this paper and the complimentary paper [11] we essentially use the framework of [18] and show how to verify this assumption in (much) more difficult situations when  $a^\alpha$  also depends on  $x$ . We emphasize that the general framework is presented in [18] and that this paper contains particular a application of that framework.

In [11] we treat the stationary case and derive error estimates for a class of problems with  $x$ -depending diffusion matrices. However, the ‘jump vector’  $\eta^\alpha$  could not depend on  $x$ . In this paper, we treat (nonlocal) time-dependent problems allowing both the diffusion matrices and the jump terms  $\eta^\alpha$  to depend on  $x$ , at least for a class of nonlinearities  $F$ . The main results are error estimates for finite difference-quadrature schemes which are compatible with the structure of  $F$ . Our work here extends the results and techniques of Krylov [21] to a nonlocal setting. It is these techniques that allow us to verify the assumptions prescribed by [18].

Throughout the major part of this paper we assume that the Lévy measure  $\nu(dz)$  sitting inside the integral operator (1.3) is bounded and compactly supported. In this case we can re-write the nonlinearity  $F$  in (1.1), possibly at the expense of changing  $b^\alpha$ , as follows

$$\begin{aligned} & \bar{F}(t, x, r, p, X, \varphi(\cdot)) \\ &= \sup_{\alpha \in \mathcal{A}} \left\{ \frac{1}{2} \operatorname{tr} [a^\alpha(t, x) X] + b^\alpha(t, x) \cdot p + \mathcal{J}^\alpha \varphi - c^\alpha(t, x) r + f^\alpha(t, x) \right\} \end{aligned}$$

where the integral operator  $\mathcal{J}^\alpha$  is defined as

$$(1.4) \quad \mathcal{J}^\alpha(\varphi)(t, x) = \int_E [\varphi(t, x + \eta^\alpha(x, z)) - \varphi(t, x)] \nu(dz).$$

Then (1.1) takes the form

$$(1.5) \quad u_t + \bar{F}(t, x, u, Du, D^2u, u(t, \cdot)) = 0 \text{ in } Q_T.$$

The general case where the Lévy measure can be unbounded and has unbounded support, can always be reduced to this case by suitable (standard) truncations. To be more precise, we replace in equation (1.1) the domain  $E$  and Lévy measure  $\nu$  by a truncated domain  $\{z : r < |z| < R\}$  and a truncated Lévy measure

$$\nu_{r,R}(dz) = \mathbf{1}_{r < |z| < R} \nu(dz).$$

Then we solve this new equation numerically using the finite-difference-quadrature method proposed in this paper. The truncation error can be controlled, and the details of this truncation procedure and its error bound can be found in [18]. In the last section of this paper we will provide a short description on the rate where the Lévy measure is singular and the cut-off is chosen optimally. Here we also give some results for the problem without truncation, but only in the case when  $\eta$  does not depend on  $x$ .

The rest of this paper is organized as follows: Section 2 collects preliminary material, including basic notations, precise form of the equations along standing assumptions on the involved coefficients, and some well-posedness and regularity results for these equations. In Section 3 we present the approximation scheme, give existence, uniqueness, comparison, and regularity results, and state our main result. Section 4 consists of the detailed proofs of the results stated in Section 3. In Section 5 we address briefly the case of unbounded Lévy measures.

## 2. PRELIMINARIES

We denote the set of all  $d \times d$  symmetric matrices  $X = (X_{ij})$ ,  $i, j = 1, 2, \dots, d$ , by  $\mathbb{S}^d$ , and let  $\mathbb{R}^d$  be the  $d$ -dimensional Euclidean space where points are denoted

by points  $x = (x^1, x^2, \dots, x^d)$ . For any  $l \in \mathbb{R}^d$ , we define the directional derivatives  $D_l$  and  $D_l^2$  as follows

$$D_l u = u_{x^i} l^i \text{ and } D_l^2 u = u_{x^i x^j} l^i l^j$$

where  $i$  and  $j$  runs from 1 to  $d$  and the summation convention applies. In this paper  $D_t$  will denote the time derivative while  $D$  will denote the spatial gradient. We denote the various constants by  $N$  or  $N(\dots)$  with or without subscripts. In the second case  $N$  only depends on the quantities in the parenthesis. Let

$$a_{\pm} = a^{\pm} = \frac{1}{2}(|a| \pm a).$$

For some set  $U$ , let  $C_b(U)$ ,  $C^2(U)$  and  $C^{1,2}(Q_T)$  denote the spaces of all functions that are bounded continuous, twice continuous differentiable, and continuous differentiable once in  $t$  and twice in  $x$  respectively. For a measurable function  $u$  defined on  $U$  we define the norm

$$|u|_0 = \text{ess sup}_{x \in U} |u(x)|.$$

For bounded functions  $u(t, x)$  and  $v(x)$  which are Lipschitz continuous in  $x$  and Hölder continuous with exponent  $\frac{1}{2}$  in  $t$ , we also define

$$|u|_{1, \frac{1}{2}} = |u|_0 + \sup_{\substack{x \neq y, t \neq s \\ x, y \in \mathbb{R}^d; t, s \in \mathbb{R}}} \frac{|u(x, t) - u(y, s)|}{|x - y| + |t - s|^{\frac{1}{2}}},$$

$$|v|_1 = |v|_0 + \sup_{x \neq y, x, y \in \mathbb{R}^n} \frac{|v(x) - v(y)|}{|x - y|}.$$

The Integro-PDE (1.5) we consider in this paper takes the following form:

$$(2.1) \quad u_t + \sup_{\alpha \in \mathcal{A}} \left\{ \mathcal{L}^{\alpha} u(t, x) + f^{\alpha}(t, x) + \mathcal{J}^{\alpha} u(t, x) \right\} = 0,$$

with the terminal condition (1.2). Here  $\mathcal{L}^{\alpha}$  is defined as

$$\mathcal{L}^{\alpha} u := a_k^{\alpha} D_{l_k}^2 u + b_k^{\alpha} D_{l_k} u - c^{\alpha} u; \quad a_k^{\alpha} := \frac{1}{2}(\sigma_k^{\alpha})^2,$$

for  $k = \pm 1, \pm 2, \dots, \pm d_1$ ,  $\mathcal{J}^{\alpha}$  is defined in (1.4), and

$$a^{\alpha}(t, x) = \frac{1}{2} \sigma^{\alpha} (\sigma^{\alpha})^T \quad \text{where} \quad \sigma_{ik}^{\alpha}(t, x) = l_k^i \sigma_k^{\alpha}(t, x).$$

Furthermore,  $l_k \in \mathbb{R}^d$ ,  $\sigma_k^{\alpha}(t, x)$ ,  $b_k^{\alpha}(t, x)$ ,  $c^{\alpha}(t, x)$  are real valued functions, and  $\eta^{\alpha}(x, z)$  is an  $\mathbb{R}^d$ -valued function.

We will assume that there are constants  $K > 1$  and  $\lambda \geq 0$  such that the following assumptions are satisfied:

- (A.1)  $\sigma_k^{\alpha}(t, x)$ ,  $\eta^{\alpha}(x, z)$ ,  $b_k^{\alpha}(t, x)$ ,  $c^{\alpha}(t, x)$ ,  $f^{\alpha}(t, x)$ ,  $u_0(x)$  are continuous in  $t, x, z, \alpha$ , and satisfy

$$l_k = -l_{-k}, \quad \sigma_k^{\alpha} = \sigma_{-k}^{\alpha}, \quad b_k^{\alpha} \geq 0, \quad c^{\alpha} \geq \lambda, \quad |l_k| \leq K.$$

- (A.2) The measure  $\nu$  is a positive Radon measure on  $E$  satisfying

$$\int_E \nu(dz) < \infty \quad \text{and} \quad \int_{E \setminus B(0, K)} \nu(dz) = 0.$$

- (A.3) For all  $\alpha, z, k$ ,  $|\sigma_k^{\alpha}|_{\frac{1}{2}, 1} + |b_k^{\alpha}|_{\frac{1}{2}, 1} + |c^{\alpha}|_{\frac{1}{2}, 1} + |f^{\alpha}|_{\frac{1}{2}, 1} + |\eta^{\alpha}(\cdot, \cdot)|_1 + |u_0|_1 \leq K$ .

Next we define the concept of viscosity solutions for (2.1).

**Definition 2.1.**  $v \in USC(Q_T)$  ( $v \in LSC(Q_T)$ ) is a viscosity *subsolution* (*supersolution*) of (2.1) if for every  $(t, x) \in Q_T$  and  $\phi \in C^{1,2}(Q_T)$  such that  $(t, x)$  is a global maximizer (global minimizer) for  $v - \phi$ ,

$$\phi_t + \sup_{\alpha \in \mathcal{A}} \left\{ a_k^\alpha D_{l_k}^2 \phi + b_k^\alpha D_{l_k} \phi - c^\alpha v(t, x) + f^\alpha(t, x) + \mathcal{J}^\alpha \phi(t, x) \right\} \geq 0 (\leq 0).$$

We say that  $v$  is a *viscosity solution* of (2.1) if  $v$  is simultaneously a sub- and supersolution of (2.1).

**Remark 2.1.** The inequalities in definition (2.1) are reversed for the sub- and supersolution compared to the usual definition (see, e.g., [16]). A time change  $t \rightarrow T - t$  will transform this terminal value problem into an (equivalent) initial value problem where the usual definition applies.

**Remark 2.2.** Contrary to the usual case [16], there are no restrictions on the growth of  $\phi$  in this definition. The reason is that the integral term is well defined whatever the growth of  $\phi$  is since the Lévy measure  $\nu$  has compact support.

For a detailed treatment for the viscosity solutions of parabolic integro-partial differential equations we suggest [16] and references therein.

In order to get the final error estimate we will use a regularizing procedure introduced by Krylov, called the method of shaking the coefficients. This procedure requires the following auxiliary equation,

$$(2.2) \quad u_t + \sup_{(\alpha, r, y) \in \mathcal{A} \times \Lambda \times B_1} \left[ \mathcal{L}^\alpha(t + \epsilon^2 r, x + \epsilon y) u + f^\alpha(t + \epsilon^2 r, x + \epsilon y) + \int_E (u(t, x + \eta^\alpha(x + \epsilon y, z)) - u(t, x)) \nu(dz) \right] = 0$$

in  $Q_T$  with the terminal data (1.2) where  $\epsilon \in \mathbb{R}$  is a constant,  $B_1 = \{x \in \mathbb{R}^d : |x| < 1\}$ , and  $\Lambda = (-1, 0)$ . We close this section by stating a well-posedness and continuous dependence result for (1.5) and (2.2). A proof can be found in [16].

**Theorem 2.1.** *Assume (A.1), (A.2), (A.3) hold, then there exist unique solutions  $v$  and  $v^\epsilon$  respectively to the terminal value problems (2.1)/(1.2) and (2.2)/(1.2) and a constant  $N$  depending only on  $d_1, K, T, \nu$  such that*

$$(2.3) \quad |v^\epsilon - v|_0 \leq N|\epsilon| \quad \text{and} \quad |v^\epsilon|_{1, \frac{1}{2}} + |v|_{1, \frac{1}{2}} \leq N.$$

*Furthermore, a comparison principle holds: If  $u$  and  $\bar{u}$  are bounded sub- and supersolutions of either (2.1)/(1.2) or (2.2)/(1.2), then  $u \leq \bar{u}$  in  $Q_T$ .*

### 3. THE DIFFERENCE-QUADRATURE SCHEME AND CONVERGENCE RATE

We begin this section with a description of a finite difference approximation to (2.1). For  $h_1, h_2, \tau > 0, l \in \mathbb{R}^d, (t, x) \in [0, T) \times \mathbb{R}^d$  we define the following finite difference operators:

$$\begin{aligned} \delta_{h_1, l} u(t, x) &= \frac{u(t, x + h_1 l) - u(t, x)}{h_1}, \\ \Delta_{h_1, l} u(t, x) &= \frac{u(t, x + h_1 l) - 2u(t, x) + u(t, x - h_1 l)}{h_1^2}, \\ \delta_\tau u(t, x) &= \frac{u(t + \tau, x) - u(t, x)}{\tau}, \end{aligned}$$

$$\delta_\tau^T u(t, x) = \frac{u(t + \tau_T(t), x) - u(t, x)}{\tau}, \quad \tau_T(t) = (t + \tau) \wedge T.$$

To discretize the integral in (2.1) we introduce a quadrature rule

$$(3.1) \quad I_{h_2}(f) = \sum_{p \in h_2 \mathbb{Z}^M} k_p f(p), \quad k_p \geq 0, \quad k_p = 0 \text{ for } |p| > K,$$

where  $p \in h_2 \mathbb{Z}^M$  and  $k_p \geq 0$  are the nodes and weights respectively. Since  $k_p \geq 0$ , this scheme is monotone. This assumption is crucial for the analysis and natural since the measure  $\nu$  is positive. Note that the sum is finite since  $k_p = 0$  for  $|p| > K$ , and this is also natural since the measure  $\nu$  has support in  $|p| \leq K$ . We also require the following consistency estimate (error estimate)

$$(3.2) \quad \left| \int_E f \nu(dz) - I_{h_2}(f) \right| \leq \nu(E) L_f h_2$$

for every Lipschitz function  $f$  with Lipschitz constant  $L_f$ .

**Remark 3.1.** Many classical quadrature rules satisfy these assumptions, the simplest example being the Riemann sum approximation,

$$I_{h_2}(f) = \sum_{p \in C h_2 \mathbb{Z}^m} f(p) \nu(p + [0, h_2]^M).$$

Other examples include the Newton-Cotes quadratures of order less than 9. We refer to [18] for a more detailed discussion.

Now we are in a position to introduce the implicit difference-quadrature scheme:

$$(3.3) \quad \delta_\tau^T u(t, x) + \sup_{\alpha \in \mathcal{A}} \left[ \mathcal{L}_{h_1}^\alpha(t, x) u + f^\alpha(t, x) + \mathcal{J}_{h_2}^\alpha u \right] = 0 \quad \text{in } Q_T,$$

with the terminal condition (1.2), where

$$\begin{aligned} \mathcal{L}_{h_1}^\alpha u &= a_k^\alpha \Delta_{h_1, l_k} u + b_k^\alpha \delta_{h_1, l_k} u - c^\alpha u, \\ \mathcal{J}_{h_2}^\alpha u &= I_{h_2}(u(t, x + \eta^\alpha(x, z)) - u(t, x)). \end{aligned}$$

As a simple consequence of Taylor's theorem, we have the following consistency bound (truncation error)

$$(3.4) \quad |\mathcal{L}_{h_1}^\alpha g(x) - \mathcal{L}^\alpha g(x)| \leq N^* \left( h_1^2 \sup_{y \in B_K(x)} |D_y^4 g| + h_1 \sup_{y \in B_K(x)} |D_y^2 g| \right),$$

for every four times differentiable function  $g$  and  $h_1 \leq 1$ , where  $N^*$  is constant which only depends on  $K, d_1$  and  $B_K(x) = \{|x| \leq K\}$ .

**Remark 3.2.** The solution  $u$  of the approximation scheme (3.3) is defined on  $Q_T$ , and not merely on a fixed grid. In part this is a technical trick to simplify the analysis, and the numerical solution defined on a grid should simply be the restriction of  $u$  to the  $h_1$ -grid. Indeed, in the local PDE context the numerical scheme would be well-defined for functions defined only on the  $h_1$ -grid. However, due to the choice of numerical quadrature, this is not the case in our nonlocal setting, and the present scheme cannot be implemented on a computer as it stands. Nevertheless, this can be remedied easily by replacing the integrand by a suitable interpolant over the  $h_1$  grid. If piecewise linear interpolation is used, monotonicity of the scheme is preserved and all the estimates obtained in this paper would still hold. From a mathematical point of view, the essential difficulties are already

present in the scheme (3.3), so to avoid increasing the length of this paper we will defer the analysis of the scheme with “interpolation” to future work.

**Remark 3.3.** The effect of using the difference operator  $\delta_\tau^T$  in (3.3) is “piecewise constant interpolation in time of the solutions”. It is equivalent to using the scheme with the operator  $\delta_\tau$  and constant-in-time initial data on the strip  $[-\tau, 0] \times \mathbb{R}^d$ .

We have the following lemma ensuring the existence of unique solution for the finite difference equation (3.3).

**Lemma 3.1.** *Assume (A.1), (A.2), (A.3), and (3.1) hold. Then there is a unique bounded function  $u(t, x)$  solving (3.3)/(1.2).*

*Proof.* For each time-level  $t$ , existence of such a solution can be proven if one know that such a solution exists for  $t + \tau_T$  by the contraction mapping argument used in the stationary case in Lemma 3.1 in [11]. Iterations, starting from terminal time  $T$  then complete the proof.  $\square$

**Remark 3.4.** It follows from the proof that the function  $u(t, x)$  is continuous in  $x$  but in general it will be discontinuous in  $t$ . However,  $u$  will satisfy a discrete Hölder bound in  $t$  (Theorem 3.4), so the size of discontinuities decrease to 0 as  $\tau \rightarrow 0$ .

For fixed  $\tau > 0$ , define

$$\bar{\mathcal{M}}_T = \{n\tau \wedge T : n = 0, 1, 2, 3, \dots\} \times \mathbb{R}^d$$

and  $\mathcal{M}_T = \bar{\mathcal{M}}_T \cap [0, T] \times \mathbb{R}^d$ . The scheme is well-defined on  $\mathcal{M}_T$ , and often we will deduce properties of the scheme on  $\bar{\mathcal{M}}_T$  and subsequently translate them to the whole space  $Q_T = \mathcal{M}_T + [0, \tau] \times \{0\}$ . We have the following lemma whose proof is postponed to the next section.

**Lemma 3.2.** *Assume (A.1), (A.2), (A.3), (3.1), (3.2) and  $h_1 < 1$ . Let  $C$  be a constant and  $u_1, u_2$  functions defined on  $\bar{\mathcal{M}}_T$ , continuous in  $x$  for each  $t$ , and for some constant  $\mu > 0$ ,*

$$\sup_{\mathcal{M}_T} |u_i(t, x) e^{-\mu|x|}| < \infty, \quad i = 1, 2.$$

If  $u_1(T, x) \leq u_2(T, x)$  and

$$(3.5) \quad \begin{aligned} & \delta_\tau^T u_1 + \sup_{\alpha \in \mathcal{A}} \left[ \mathcal{L}_{h_1}^\alpha u_1 + f^\alpha(t, x) + \mathcal{J}_{h_2}^\alpha u_1 \right] + C \\ & \geq \delta_\tau^T u_2 + \sup_{\alpha \in \mathcal{A}} \left[ \mathcal{L}_{h_1}^\alpha u_2 + f^\alpha(t, x) + \mathcal{J}_{h_2}^\alpha u_2 \right], \end{aligned}$$

then there exists a constant  $\tau^* > 0$  depending only on  $K, d_1, \mu, \nu(E)$  such that if  $\tau \in (0, \tau^*)$ ,

$$(3.6) \quad u_1 \leq u_2 + (T + \tau)C_+ \quad \text{in } \bar{\mathcal{M}}_T.$$

Furthermore,  $\tau^*(K, d_1, \mu, \nu(E)) \rightarrow \infty$  as  $\mu \downarrow 0$ , and if  $u_1, u_2$  are bounded, (3.6) holds for all  $\tau > 0$ .

**Corollary 3.3.** *Assume (A.1), (A.2), (A.3), (3.1), (3.2) and  $h_1 < 1$ . Then the solution  $v_{\tau, h}$  of (3.3)-(1.2) satisfies*

$$|v_{\tau, h}|_0 \leq K(T + \tau) + |u_0|_0.$$

*Proof.* The function  $\pm[K(T - t) + |u_0|_0]$  is supersolution/subsolution of (3.3)-(1.2) (remember  $c^\alpha \geq \lambda \geq 0$ ), so the result follows from Lemma 3.4.  $\square$

Consider the terminal value problem

$$(3.7) \quad \delta_\tau^T u + \sup_{(\alpha, r, y) \in A \times \Lambda \times B_1} \left[ \mathcal{L}_{h_1}^\alpha(t + \epsilon^2 r, x + \epsilon y) u(t, x) + f^\alpha(t + \epsilon^2 r, x + \epsilon y) + \sum_p k_p (u(t, x + \eta^\alpha(x + \epsilon y, p)) - u(t, x)) \right] = 0 \text{ in } Q_T,$$

with terminal data (1.2). This is the difference-quadrature scheme corresponding to (2.2). By Lemma 3.1 and Corollary 3.3 there exists a unique bounded solution  $v_{\tau, h}^\epsilon$  of this problem.

We have the following theorem, whos proof will be given in the next section.

**Theorem 3.4.** *Assume (A.1), (A.2), (A.3), (3.1), (3.2),  $0 \leq h_1 \leq 1$ , and  $0 \leq \tau \leq \tau_0$ . If  $\tau_0$  is small enough, there exists a constant  $N$  (depending only on  $\tau_0, \lambda, T, K, d_1$ , and  $\nu(E)$ ); such that for all  $\epsilon \in \mathbb{R}$*

$$(3.8) \quad |v_{\tau, h}^\epsilon(t, x) - v_{\tau, h}(t, x)| \leq N|\epsilon|,$$

$$(3.9) \quad |v_{\tau, h}^\epsilon(t, x) - v_{\tau, h}^\epsilon(t, y)| + |v_{\tau, h}(t, x) - v_{\tau, h}(t, y)| \leq N|x - y|,$$

$$(3.10) \quad |v_{\tau, h}^\epsilon(t, x) - v_{\tau, h}^\epsilon(s, x)| + |v_{\tau, h}(t, x) - v_{\tau, h}(s, x)| \leq N(|t - s|^{\frac{1}{2}} + \tau^{\frac{1}{2}}),$$

for all  $(t, x), (s, y) \in \bar{Q}_T$ .

Now, with the help of the results stated above, we are in a position to prove the main contribution of this paper, namely

**Theorem 3.5.** *Assume (A.1), (A.2), (A.3), (3.1), (3.2),  $0 < h_1, h_2$ , and  $0 < \tau \leq \tau_0$ . If  $\tau_0, h_1, h_2$  are small enough, there exists a constant  $N_1$  (depending only on  $\tau_0, \lambda, d_1, T, K, \nu(E)$ ), such that*

$$|v - v_{\tau, h}|_0 \leq N_1 [\tau^{\frac{1}{4}} + h_1^{\frac{1}{2}} + h_2].$$

*Proof.* Take  $\epsilon = (\tau + h_1^2 + h_2^4)^{\frac{1}{4}}$  and let  $\tau_0, h_1, h_2$  be sufficiently small such that  $\epsilon < 1$ . If  $T < 2\epsilon^2$  then the theorem holds because by (3.10) and (2.3) and the definition of  $\epsilon$ ,

$$\begin{aligned} \sup_{Q_T} |v_{\tau, h} - v| &\leq \sup_{Q_T} |v_{\tau, h} - u_0| + \sup_{Q_T} |u_0 - v| \\ &\leq N(T^{\frac{1}{2}} + \tau^{\frac{1}{2}}) \leq N(\tau + h_1^2 + h_2^4)^{\frac{1}{4}}. \end{aligned}$$

Next we consider the case  $T > 2\epsilon^2$ . First we prove the upper bound

$$(3.11) \quad v - v_{\tau, h} \leq N(\tau^{\frac{1}{4}} + h_1^{\frac{1}{2}} + h_2).$$

For each  $\alpha \in \mathcal{A}, r \in (-1, 0)$  and  $|y| < 1$ , equation (3.7) implies

$$(3.12) \quad \delta_\tau^T v_{\tau, h}^\epsilon(t - \epsilon^2 r, x - \epsilon y) + \mathcal{L}_{h_1}^\alpha(t, x) v_{\tau, h}(t - \epsilon^2 r, x - \epsilon y) + f^\alpha(t, x) + \sum_p k_p \left[ v_{\tau, h}^\epsilon(t - \epsilon^2 r, x + \eta^\alpha(x, p) - \epsilon y) - v_{\tau, h}^\epsilon(t - \epsilon^2 r, x - \epsilon y) \right] \leq 0$$

for  $(t, x) \in \bar{Q}_{T - \epsilon^2}$ .

Now use Krylov's technique i.e. multiply inequality (3.12) with a mollifier and convolve. Let  $\zeta \in C_0^\infty(\mathbb{R}^{d+1})$  be our mollifier, a positive function with unit integral and having support in  $\Lambda \times B_1$ . Also denote,

$$u^{(\epsilon)}(t, x) = \epsilon^{(-d-2)} \int_{\mathbb{R}^{d+1}} u(t - s, x - y) \zeta\left(\frac{s}{\epsilon^2}, \frac{y}{\epsilon}\right) ds dy$$

Then multiplying (3.12) by  $\epsilon^{-d-2}\zeta(s/\epsilon^2, y/\epsilon)$  and integrating with respect to  $(s, y)$  we obtain, for each  $\alpha \in \mathcal{A}$ , on  $\bar{Q}_{T-2\epsilon^2}$

$$\delta_\tau^T v_{\tau,h}^{\epsilon(\epsilon)} + \mathcal{L}_{h_1}^\alpha(t, x)v_{\tau,h}^{\epsilon(\epsilon)} + f^\alpha + \sum_p k_p(v_{\tau,h}^{\epsilon(\epsilon)}(t, x + \eta^\alpha(x, p)) - v_{\tau,h}^{\epsilon(\epsilon)}) \leq 0.$$

From (3.4), (3.2), and Taylor's formula we have

$$\begin{aligned} & \frac{\partial}{\partial t} v_{\tau,h}^{\epsilon(\epsilon)} + \mathcal{L}^\alpha(t, x)v_{\tau,h}^{\epsilon(\epsilon)} + f^\alpha + \int_{\mathbb{R}^M \setminus \{0\}} \left( v_{\tau,h}^{\epsilon(\epsilon)}(t, x + \eta^\alpha(x, z)) - v_{\tau,h}^{\epsilon(\epsilon)} \right) \nu(dz) \\ & \leq N \left( \tau |D_t^2 v_{\tau,h}^{\epsilon(\epsilon)}|_{0, \bar{Q}_{T-2\epsilon^2}} + h_1^2 |D_x^4 v_{\tau,h}^{\epsilon(\epsilon)}|_{0, \bar{Q}_{T-2\epsilon^2}} + h_1 |D_x^2 v_{\tau,h}^{\epsilon(\epsilon)}|_{0, \bar{Q}_{T-2\epsilon^2}} \right. \\ & \quad \left. + h_2 |D_x v_{\tau,h}^{\epsilon(\epsilon)}|_{0, \bar{Q}_{T-2\epsilon^2}} \right) := I \quad \text{in } \bar{Q}_{T-2\epsilon^2}. \end{aligned}$$

Clearly  $v_{\tau,h}^{\epsilon(\epsilon)} + (T - 2\epsilon^2 - t)I$  is a classical supersolution to the equation (2.1) and hence a viscosity supersolution as well in  $Q_{T-2\epsilon^2}$ . Now using the comparison principle (Theorem 2.1) we have

$$(3.13) \quad v \leq v_{\tau,h}^{\epsilon(\epsilon)} + (T - 2\epsilon^2 - t)I + \sup_{\{(T-2\epsilon^2) \times \mathbb{R}^d\}} |v - v^{\epsilon(\epsilon)}|.$$

Using properties of convolutions and regularity of  $v_{\tau,h}^\epsilon$  (Theorem 3.4),

$$|v_{\tau,h}^{\epsilon(\epsilon)} - v_{\tau,h}^\epsilon| \leq N\epsilon \quad \text{and} \quad \epsilon^{2n-1} |D_t^n v_{\tau,h}^{\epsilon(\epsilon)}|_{0, \bar{Q}_{T-2\epsilon^2}} + \epsilon^{n-1} |D_x^n v_{\tau,h}^{\epsilon(\epsilon)}|_{0, \bar{Q}_{T-2\epsilon^2}} \leq N.$$

By the same reasoning as in the beginning of the proof, we also find

$$|v(T - 2\epsilon^2, x) - v^{\epsilon(\epsilon)}(T - 2\epsilon^2, x)| \leq N\epsilon.$$

By the above estimates, Theorem 3.4, and recalling that  $\epsilon^4 = \tau + h_1^2 + h_2^4$ , we obtain

$$\begin{aligned} v & \leq v_{\tau,h}^\epsilon + N \left( \epsilon + \frac{\tau + h_1^2}{\epsilon^3} + \frac{h_1}{\epsilon} + h_2 \right) \\ & \leq v_{\tau,h} + N \left( \epsilon + \frac{\tau + h_1^2}{\epsilon^3} + \frac{h_1}{\epsilon} + h_2 \right) \\ & \leq v_{\tau,h} + N(\tau + h_1^2 + h_2^4)^{\frac{1}{4}} \quad \text{in } \bar{Q}_{T-2\epsilon^2}. \end{aligned}$$

By the regularity of  $v, v_{\tau,h}$  and the argument given in the case  $T < 2\epsilon^2$ , this estimate in fact holds in all of  $Q_T$ . Moreover, it can be checked that all constants  $N$  only depend on  $\tau_0, \lambda, \nu(E), d_1, d, K$  and  $T$ . This completes the proof of (3.11).

The lower bound

$$(3.14) \quad v_{\tau,h} - v \leq N(\tau^{\frac{1}{4}} + h_1^{\frac{1}{2}} + h_2),$$

can be proved in a similar way. Interchange the role of the finite difference scheme and the equation (2.1) in the argument leading to (3.11). Now it can be shown that  $v^{\epsilon(\epsilon)}$  is a classical supersolution of (2.1) in  $Q_{T-\epsilon^2}$ . We skip the arguments since they are similar to the arguments for stationary integro-PDEs given in [18], see also [20, 15] for time-dependent pure PDEs.

By consistency (3.2) and (3.4), regularity of  $v^{\epsilon(\epsilon)}$ , and properties of mollifiers, it follows that

$$\delta_\tau^T v^{\epsilon(\epsilon)} + \sup_{\alpha \in \mathcal{A}} [L_{h_1}^\alpha v^{\epsilon(\epsilon)} + f^\alpha + \mathcal{J}_{h_2}^\alpha v^{\epsilon(\epsilon)}] \leq N \left( \frac{\tau + h_1^2}{\epsilon^3} + \frac{h_1}{\epsilon} + h_2 \right)$$

in  $\bar{Q}_{T-\epsilon^2-\tau}$ , and the comparison result (Lemma 3.2) then gives

$$(3.15) \quad v_{\tau,h} \leq v^{\epsilon(\epsilon)} + \sup_{Q_T \setminus Q_{T-\epsilon^2-\tau}} (v_{\tau,h} - v^{\epsilon(\epsilon)})_+ + N \left( \frac{\tau + h_1^2}{\epsilon^3} + \frac{h_1}{\epsilon} + h_2 \right).$$

Since  $\tau \leq \epsilon^4 \leq \epsilon^2 \leq 1$  and hence  $Q_T \setminus Q_{T-\epsilon^2-\tau} \subset Q_T \setminus Q_{T-2\epsilon^2}$ ,

$$\sup_{Q_T \setminus Q_{T-\epsilon^2-\tau}} (v_{\tau,h} - v^{\epsilon(\epsilon)})_+ \leq \sup_{Q_T \setminus Q_{T-2\epsilon^2}} |v_{\tau,h} - v^{\epsilon(\epsilon)}| \leq N\epsilon,$$

where the last inequality was proved at the start of this proof. This estimate, (3.15), and the definition of  $\epsilon$ , implies the lower bound (3.14). It can be checked that  $N$  depends only on  $\nu(E)$ ,  $K$ ,  $d_1$ ,  $d$  and  $T$ .  $\square$

#### 4. PROOFS OF THE RESULTS STATED IN SECTION 3

In this section we prove comparison and Lipschitz continuity results for the solution of the difference-quadrature scheme (3.3), (1.2). As an application of the Lipschitz result we derive a continuous dependence estimate for the scheme. Although the basic ideas behind our proofs come from Krylov [21], the nonlocal nature of the problem adds to some extra difficulties and they do not allow us to adopt the ‘‘local’’ approach of Krylov. Our approach is more direct and we employ some new techniques.

We begin by stating some auxiliary results. To this end, we need the translation operator

$$T_{h_1,l}u(x) := u(x + h_1l).$$

We now give some technical lemmas whose proofs can be found in [21].

**Lemma 4.1.** *For any functions  $u(x), v(x), h_1 > 0$ . and  $l \in \mathbb{R}^d$  we have*

$$\begin{aligned} T_{h_1,-l}T_{h_1,l}u &= u, \\ T_{h_1,l}\delta_{h_1,-l} &= \delta_{h_1,-l}T_{h_1,l} = -T_{h_1,-l}\delta_{h_1,l} = -\delta_{h_1,l}T_{h_1,-l} = -\delta_{h_1,-l}, \\ \delta_{h_1,l}(uv) &= v\delta_{h_1,l}(u) + T_{h_1,l}u\delta_{h_1,l}(v), \\ &= u\delta_{h_1,l}(v) + v\delta_{h_1,l}(u) + h_1(\delta_{h_1,l}(v))(\delta_{h_1,l}(u)) \\ \Delta_{h_1,l}(uv) &= u\Delta_{h_1,l}(v) + v\Delta_{h_1,l}(u) + (\delta_{h_1,l}(v))(\delta_{h_1,l}(u)) \\ &\quad + (\delta_{h_1,-l}(v))(\delta_{h_1,-l}(u)). \end{aligned}$$

*In particular,*

$$\Delta_{h_1,l}(u^2) = 2u\Delta_{h_1,l}u + (\delta_{h_1,l}u)^2 + (\delta_{h_1,-l}u)^2.$$

**Lemma 4.2.** *Let  $u, v, w$  be functions on  $\mathbb{R}^d, l, x_0 \in \mathbb{R}^d, h_1 > 0$ . Assume that  $v(x_0) \leq 0$ . Then at  $x_0$  it holds*

$$(4.1) \quad -\delta_{h_1,l}v \leq \delta_{h_1,l}(v_-), \quad -\Delta_{h_1,l}v \leq \Delta_{h_1,l}(v_-),$$

$$(4.2) \quad |\Delta_{h_1,l}u| \leq |\delta_{h_1,-l}((\delta_{h_1,l}u)_-)| + |\delta_{h_1,l}((\delta_{h_1,-l}u)_-)|,$$

$$|\Delta_{h_1,l}u| \leq |\delta_{h_1,-l}((\delta_{h_1,l}u)_+)| + |\delta_{h_1,l}((\delta_{h_1,-l}u)_+)|.$$

Now we prove Lemma 3.2.

*Proof of Lemma 3.2:* Let  $T'$  be the smallest  $j\tau$  which exceeds  $T$ , where  $j \in \mathbb{N}$ . A solution to the equation (3.3) on  $\bar{\mathcal{M}}_T$  could be viewed as a solution to the same on  $\bar{\mathcal{M}}_{T'}$  after trivially redefining the function on  $\{T'\} \times \mathbb{R}^d$ . So without loss of generality we assume that  $T = T'$ .

From (3.5) we then have in  $\mathcal{M}_T$ , for  $u = u_1 - u_2$

$$\delta_\tau u + \sup_{\alpha \in \mathcal{A}} \left[ \mathcal{L}_{h_1}^\alpha u + \sum_{p \in h_2 \mathbb{Z}^M} k_p(u(t, x + \eta^\alpha(x, p)) - u(t, x)) \right] + C \geq 0.$$

Let  $w = u - C_+(T - t)$  and note that

$$\begin{aligned} & \delta_\tau w + \sup_{\alpha \in \mathcal{A}} \left[ \mathcal{L}_{h_1}^\alpha w + \sum_{p \in h_2 \mathbb{Z}^M} k_p(w(t, x + \eta^\alpha(x, p)) - w(t, x)) \right] \\ & \geq \delta_\tau u + \sup_{\alpha \in \mathcal{A}} \left[ \mathcal{L}_{h_1}^\alpha u + \sum_{p \in h_2 \mathbb{Z}^M} k_p(u(t, x + \eta^\alpha(x, p)) - u(t, x)) \right] + C_+ + \lambda C_+(T - t) \\ & \geq 0, \end{aligned}$$

and hence for  $\varepsilon > 0$ ,

$$w + \varepsilon \delta_\tau w + \varepsilon \sup_{\alpha \in \mathcal{A}} \left[ \mathcal{L}_{h_1}^\alpha w + \sum_{p \in h_2 \mathbb{Z}^M} k_p(w(t, x + \eta^\alpha(x, p)) - w(t, x)) \right] \geq w.$$

For any  $\psi \geq w$ , we can choose  $\varepsilon$  small enough so that in  $\mathcal{M}_T$ ,

$$\begin{aligned} & \psi + \varepsilon \delta_\tau \psi + \varepsilon \sup_{\alpha \in \mathcal{A}} \left[ \mathcal{L}_{h_1}^\alpha \psi + \sum_{p \in h_2 \mathbb{Z}^M} k_p(\psi(t, x + \eta^\alpha(x, p)) - \psi(t, x)) \right] \\ (4.3) \quad & \geq w + \varepsilon \delta_\tau w + \varepsilon \sup_{\alpha \in \mathcal{A}} \left[ \mathcal{L}_{h_1}^\alpha w + \sum_{p \in h_2 \mathbb{Z}^M} k_p(w(t, x + \eta^\alpha(x, p)) - w(t, x)) \right] \geq w. \end{aligned}$$

For a constant  $\gamma$  and  $\langle x \rangle := \sqrt{1 + x^2}$ , we define the functions  $\xi(t)$ ,  $\beta(x)$ , and  $\zeta(t, x)$  on  $\bar{\mathcal{M}}_T$  in the following way:

$$\begin{aligned} \xi(T) &= 1, \quad \xi(t) = \gamma^{-1} \xi(t + \tau_T(t)) \text{ if } t \in [0, T) \\ \beta(x) &= \cosh(\mu \langle x \rangle), \quad \zeta(t, x) = \xi(t) \beta(x). \end{aligned}$$

Note that  $\xi$  is recursively defined. By straightforward computations we have,

$$\begin{aligned} & \sup_{\alpha \in \mathcal{A}} \left[ \mathcal{L}_{h_1}^\alpha \beta + \sum_{p \in h_2 \mathbb{Z}^M} k_p(\beta(x + \eta^\alpha(x, p)) - \beta(x)) \right] \\ & \leq \sup_{\alpha \in \mathcal{A}} \mathcal{L}^\alpha \beta + N_1(h_1^2 + h_1) \cosh(\mu \langle x \rangle + K) + N_2(\nu(E), \mu) \cosh(\mu \langle x \rangle + K) \\ & \leq N_2 \cosh(\mu \langle x \rangle + K). \end{aligned}$$

Since  $\delta_\tau \xi(t) = \frac{\gamma-1}{\tau} \xi(t)$  and  $\cosh(\mu \langle x \rangle + K) \leq e^K \cosh(\mu \langle x \rangle)$ , it follows that

$$\begin{aligned} & \delta_\tau \zeta + \sup_{\alpha \in \mathcal{A}} \left[ \mathcal{L}_{h_1}^\alpha \zeta + \sum_{p \in h_2 \mathbb{Z}^M} k_p(\zeta(t, x + \eta^\alpha(x, p)) - \zeta(t, x)) \right] \\ & \leq \tau^{-1}(\gamma - 1)\zeta + N_3 \zeta = \kappa(\gamma)\zeta, \end{aligned}$$

where  $N_3 = N_2 e^K$  and  $\kappa(\gamma) = \tau^{-1}(\gamma - 1) + N_3$ . We take  $\tau_* = N_3^{-1}$  and let  $\tau < \tau_*$ . Then  $\kappa(0) < 0$  and  $\kappa(1) \geq 0$  and hence we can choose  $\gamma$  so that  $\kappa < 0$  and  $1 + \kappa \varepsilon > 0$  for all  $\varepsilon$  small enough.

Now set  $N = \sup_{\mathcal{M}_T} \frac{w_\pm}{\zeta}$ . Taking  $\psi = N\zeta$  and  $\varepsilon$  small enough, (4.3) leads to

$$\begin{aligned} & N\zeta(1 + \kappa \varepsilon) = N\zeta + \varepsilon N\delta_\tau \zeta \\ & \geq \psi + \varepsilon \delta_\tau \psi + \varepsilon \sup_{\alpha \in \mathcal{A}} \left[ \mathcal{L}_{h_1}^\alpha \psi + \sum_{p \in h_2 \mathbb{Z}^M} k_p(\psi(t, x + \eta^\alpha(x, p)) - \psi(t, x)) \right] \geq w. \end{aligned}$$

in  $\mathcal{M}_T$ . But  $w(T, x)$  is negative by definition, so the inequality holds on entire  $\bar{\mathcal{M}}_T$ . By the definition of  $N$ , we then have  $N(1 + \kappa\epsilon) \geq N$ . Since  $\kappa < 0$ , we conclude that  $N = 0$  and hence  $w \leq 0$  and (3.6) follows. The remaining part of the lemma becomes obvious if we choose  $N_1 = N_2 = N_3 = 0$   $\square$

Next we state and prove the key technical result of this paper.

**Theorem 4.3.** *Assume (A.1), (A.3), (A.2), (3.1) and (3.2) hold. Let  $u(t, x)$  be a function on  $\bar{\mathcal{M}}_T$  solving (3.3) with  $|u(T, \cdot)|_1 < \infty$ . There is a constant  $N > 0$ , depending only on  $K, d_1, d$  and the Lévy measure  $\nu$ , such that, if there is a number  $c_0 \geq 0$  satisfying*

$$(4.4) \quad \lambda + \frac{1 - e^{-c_0\tau}}{\tau} > N,$$

then for every  $0 < \epsilon < Kh_1, l \in \mathbb{R}^d$ ,

$$|\delta_{\epsilon, \pm l} u(t, x)| \leq N_1(1 \vee |l|) \left( 1 + \sup_{k, x} |\delta_{h_1, l_k} u(T, \cdot)| + \sup_x |\delta_{\epsilon, \pm l} u(T, \cdot)| \right) \quad \text{in } \bar{\mathcal{M}}_T,$$

where  $N_1$  only depend on  $T, \lambda, c_0, K, d, d_1, \nu(E)$ .

*Proof.* We start by introducing a few additional notations. Let  $r$  and  $k$  be indices running through  $\{\pm 1, \pm 2, \dots, \pm(d_1 + 1)\}$  and  $\{\pm 1, \pm 2, \dots, \pm d_1\}$  respectively, let  $0 < \epsilon \leq Kh_1$ , and define

$$h_k = h_1, \quad k = \pm 1, \pm 2, \dots, \pm d_1, \quad h_{\pm(d_1+1)} = \epsilon, \quad l_{\pm(d_1+1)} = \pm l.$$

Choose a constant  $c_0 \geq 0$ , let  $T'$  be the least  $n\tau$ ,  $n = 1, 2, 3, \dots$ , such that  $n\tau \geq T$ , and define

$$\begin{aligned} \xi(t) &= e^{c_0 t} \quad \text{if } t < T', \quad \xi(T) = e^{c_0 T'} \quad \text{otherwise;} \\ v &= \xi u; \\ v_r &= \delta_{h_r, l_r} v \quad \text{if } r = \pm 1, \pm 2, \dots, \pm d_1; \\ v_{\pm(d_1+1)} &= \frac{v(t, x \pm \epsilon l) - v(t, x)}{\epsilon(1 \vee |l|)}; \\ M &= \sup_{(t, x) \in \bar{Q}_T} |v(t, x)|, \quad M_1 = \sup_{r, x, l, t} |v_r|. \end{aligned}$$

Now define

$$W(t, x, l) = \sum_r (v_r^-)^2 \quad \text{and} \quad V(t, x, l) = W(t, x, l) - \delta C(x),$$

where  $\delta > 0$  and  $C(x) \in C^2(\mathbb{R}^d)$  is positive, convex, and satisfy

$$\lim_{|x| \rightarrow \infty} C(x) = \infty \quad \text{and} \quad |DC|_0 + |D^2C|_0 < \infty.$$

To prove the theorem we have to find a bound on  $M_1$  which is independent of the discretization constants. We will derive such a bound for  $W$ , and towards the end of the proof we will show that this bound implies the sought after bound on  $M_1$ .

From the properties of  $C(x)$ , it is clear that  $V(t, x, l)$  is bounded above and that there exists a point  $(t_0, x_0, l_0) \in \bar{\mathcal{M}}_T \times \mathbb{R}^d$  such that

$$V(t_0, x_0, l_0) = \sup_{(t, x, l)} V(t, x, l).$$

If  $t_0 = T$ , then

$$(4.5) \quad V(T, \cdot) \leq W(T, \cdot) \leq N(d_1)e^{2c_0T'} \left( \sup_{r,x} u_r(T, x) \right)^2,$$

and the theorem is true by Lipschitz continuity of  $u(T, x)$ .

From now on we take  $t_0 < T$ . By the definition of supremum, there is a sequence of control parameters  $(\alpha_n) \in \mathcal{A}$  (depending on the maximum point  $(x_0, t_0, l_0)$ ) such that

$$\begin{aligned} & \lim_{n \rightarrow \infty} \left[ \mathcal{L}_{h_1}^{\alpha_n}(t_0, x_0)u(t_0, x_0) + f^{\alpha_n}(t_0, x_0) + \mathcal{J}_{h_2}^{\alpha_n}u(t_0, x_0) \right] \\ &= \sup_{\alpha \in \mathcal{A}} \left[ \mathcal{L}_{h_1}^{\alpha}(t_0, x_0)u(t_0, x_0) + f^{\alpha}(t_0, x_0) + \mathcal{J}_{h_2}^{\alpha}(u(t_0, x_0)) \right]. \end{aligned}$$

By assumption (A.3) and the Arzela-Ascoli theorem there is a subsequence  $\{\alpha_n\}$  and functions  $\bar{a}_k, \bar{b}_k, \bar{c}, \bar{f}, \bar{\eta}$ , such that

$$(a_k^{\alpha_n}, b_k^{\alpha_n}, c^{\alpha_n}, f^{\alpha_n}, \eta^{\alpha_n}) \rightarrow (\bar{a}_k, \bar{b}_k, \bar{c}, \bar{f}, \bar{\eta}) \quad \text{locally uniformly.}$$

Obviously,  $\bar{a}_k, \bar{b}_k, \bar{c}, \bar{f}, \bar{\eta}$  also satisfy (A.1) and (A.3). Moreover since  $u$  solve (3.3),

$$(4.6) \quad \begin{aligned} & \delta_\tau^T u + \bar{a}_k \Delta_{h_1, l_k} u + \bar{b}_k \delta_{h_1, l_k} u - \bar{c}u + \bar{f} \\ & + \sum_{p \in h_2 \mathbb{Z}^M} k_p (u(t_0, x_0 + \bar{\eta}(x_0, p)) - u(t_0, x_0)) = 0, \end{aligned}$$

at the point  $(t_0, x_0)$ , while at the points  $(t_0, x_0 + h_r l_r)$ ,

$$(4.7) \quad \begin{aligned} & \delta_\tau^T u + \bar{a}_k \Delta_{h_1, l_k} u + \bar{b}_k \delta_{h_1, l_k} u - \bar{c}u + \bar{f} \\ & + \sum_{p \in h_2 \mathbb{Z}^M} k_p (u(\cdot, \cdot + \bar{\eta}(\cdot, p)) - u(\cdot, \cdot)) \Big|_{(t_0, x_0 + h_r l_r)} \leq 0. \end{aligned}$$

The last inequality holds at every point in  $Q_T$ . For simplicity we now drop the 0 subscript and rename the maximum point  $(x, t, l)$ . Replacing  $u$  by  $\xi^{-1}v$  in (4.6) and (4.7) we get

$$(4.8) \quad \begin{aligned} & \delta_\tau^T (\xi^{-1}v) + \xi^{-1} \left( \bar{a}_k \Delta_{h_1, l_k} v + \bar{b}_k \delta_{h_1, l_k} v - \bar{c}v + \bar{f} \right. \\ & \left. + \sum_{p \in h_2 \mathbb{Z}^M} k_p (v(t, x + \bar{\eta}(x, p)) - v(t, x)) \right) = 0 \end{aligned}$$

at the point  $(t, x)$  and for each  $r$ , and at the points  $(t, x + h_r l_r)$  we have

$$(4.9) \quad \begin{aligned} & \left[ \delta_\tau^T (\xi^{-1}v) + \xi^{-1} \left( \bar{a}_k \Delta_{h_1, l_k} v + \bar{b}_k \delta_{h_1, l_k} v - \bar{c}v + \bar{f} \right. \right. \\ & \left. \left. + \sum_{p \in h_2 \mathbb{Z}^M} k_p (v(\cdot, \cdot + \bar{\eta}(\cdot, p)) - v(\cdot, \cdot)) \right) \right] \Big|_{(t, x + h_r l_r)} \leq 0. \end{aligned}$$

Subtracting (4.8) from (4.9) and dividing the result by  $h_r$ , for  $r = \pm 1, \pm 2, \dots, \pm d_1$ , and by  $\epsilon(|l| \vee 1)$  for  $r = \pm(d_1 + 1)$ , and using the product rule for difference quotients (Lemma 4.1) we get

$$(4.10) \quad \delta_\tau^T (\xi^{-1}v_r) + \xi^{-1} \left[ \bar{a}_k \Delta_{h_k, l_k} v_r + I_{1r} + I_{2r} + I_{3r} + I_{4r} + I_{5r} \right] \leq 0,$$

where there is no summation with respect to  $r$ . Here

$$I_{1r} = \begin{cases} (\delta_{h_r, l_r} \bar{a}_k) \Delta_{h_k, l_k} v, & \text{if } r \neq \pm(d_1 + 1) \\ \frac{1}{(1V|l|)} (\delta_{h_r, l_r} \bar{a}_k) \Delta_{h_k, l_k} v, & \text{if } r = \pm(d_1 + 1), \end{cases}$$

$$\begin{aligned}
I_{2r} &= h_r (\delta_{h_r, l_r} \bar{a}_k) \Delta_{h_k, l_k} v_r, \\
I_{3r} &= \begin{cases} (T_{h_r, l_r} \bar{b}_k) \delta_{h_k, l_k} v_r + (\delta_{h_r, l_r} \bar{b}_k) \delta_{h_k, l_k} v, & \text{if } r \neq \pm(d_1 + 1) \\ (T_{h_r, l_r} \bar{b}_k) \delta_{h_k, l_k} v_r + \frac{1}{(1 \vee |l|)} (\delta_{h_r, l_r} \bar{b}_k) \delta_{h_k, l_k} v, & \text{if } r = \pm(d_1 + 1), \end{cases} \\
I_{4r} &= \begin{cases} -(\delta_{h_r, l_r} \bar{c}) v - (T_{h_r, l_r} \bar{c}) v_r + \xi \delta_{h_r, l_r} \bar{f}, & \text{if } r \neq \pm(d_1 + 1) \\ -\frac{1}{(1 \vee |l|)} (\delta_{h_r, l_r} \bar{c}) v - (T_{h_r, l_r} \bar{c}) v_r + \frac{1}{(1 \vee |l|)} \xi \delta_{h_r, l_r} \bar{f}, & \text{if } r = \pm(d_1 + 1), \end{cases} \\
I_{5r} &= \begin{cases} \sum_p k_p \frac{(v(t, x + h_r l_r + \bar{\eta}(x + h_r l_r, p)) - v(t, x + \bar{\eta}(x, p)))}{h_r} - \nu(E) v_r, & \text{if } r \neq \pm(d_1 + 1) \\ \sum_p k_p \frac{(v(t, x + h_r l_r + \bar{\eta}(x + h_r l_r, p)) - v(t, x + \bar{\eta}(x, p)))}{h_r (1 \vee |l|)} - \nu(E) v_r, & \text{if } r = \pm(d_1 + 1). \end{cases}
\end{aligned}$$

The last term is of particular relevance to this paper as it comes from the discretization of the integral term.

Now multiply (4.10) by  $\xi v_r^-$  and sum up with respect to  $r$ . The main part of the proof involves the estimation of each of the above terms as they appear after summation had been done.

We start with the term  $\sum_r v_r^- I_{5r}$ . Note that  $v_r^- v_r = (v_r^-)^2$  and moreover that  $\sum_p k_p = \nu(E)$  by (3.2). We get

$$\begin{aligned}
& \sum_r v_r^- I_{5r} \\
&= \nu(E) W + \sum_{p, r \neq \pm(d_1 + 1)} k_p v_r^- \frac{v(t, x + h_r l_r + \bar{\eta}(x + h_r l_r, p)) - v(t, x + \bar{\eta}(x, p))}{h_r} \\
&+ \sum_{p, r = \pm(d_1 + 1)} k_p v_r^- \frac{v(t, x + h_r l_r + \bar{\eta}(x + h_r l_r, p)) - v(t, x + \bar{\eta}(x, p))}{h_r (1 \vee |l_r|)}.
\end{aligned}$$

For  $r = \pm 1, \pm 2, \dots, \pm d_1$  we have

$$\begin{aligned}
& \left| \frac{v(t, x + h_r l_r + \bar{\eta}(x + h_r l_r, p)) - v(t, x + \bar{\eta}(x, p))}{h_r} \right| \\
&= \left( \frac{\epsilon}{h_r} \right) (| \frac{h_r}{\epsilon} l' | \vee 1) \left| \frac{v(t, x + \bar{\eta}(x, p) + \epsilon \frac{h_r l'}{\epsilon}) - v(t, x + \bar{\eta}(x, p))}{\epsilon (1 \vee | \frac{h_r}{\epsilon} l' |)} \right| \\
&\leq K |l'| M_1,
\end{aligned}$$

since  $\epsilon \leq K h_1$  and where

$$l' = l_r + \frac{\bar{\eta}(x + h_r l_r, p) - \bar{\eta}(x, p)}{h_r} \quad \text{and hence} \quad |l'| \leq |l_r| + |l_r| |\nabla \bar{\eta}(\cdot, p)|_{L^\infty}.$$

For  $r = \pm(d_1 + 1)$  similarly we have,

$$\left| \frac{v(t, x + \epsilon l + \bar{\eta}(x + \epsilon l, p)) - v(t, x + \bar{\eta}(x, p))}{\epsilon (1 \vee |l|)} \right| \leq M_1 \frac{1 \vee (|l'|)}{1 \vee |l|}$$

where

$$l' = l + \frac{\bar{\eta}(x + \epsilon l, p) - \bar{\eta}(x, p)}{\epsilon} \quad \text{and hence} \quad |l'| \leq |l| (1 + |\nabla \bar{\eta}(\cdot, p)|_{L^\infty}).$$

Putting the above pieces together and using Cauchy-Schwartz inequality we get,

$$(4.11) \quad \sum_r v_r^- I_{5r} \geq \nu(E) W - N(d_1, K) \nu(E) M_1^2.$$

Next, we estimate the term  $\sum_r v_r^- I_{3r}$ :

$$(4.12) \quad \begin{aligned} \sum_r v_r^- I_{3r} &= \sum_r v_r^- T_{h_r, l_r} \bar{b}_k \delta_{h_k, l_k} v_r + \sum_{r \neq \pm(d_1+1)} v_r^- (\delta_{h_r, l_r} \bar{b}_k) \delta_{h_k, l_k} v \\ &\quad + \sum_{r=\pm(d_1+1)} \frac{1}{1 \vee |l|} v_r^- (\delta_{h_r, l_r} \bar{b}_k) \delta_{h_k, l_k} v. \end{aligned}$$

At the maximum point  $(t, x, l)$  for  $V$ , Lemma 4.1 and (4.1) yields for each  $k$

$$0 \geq \delta_{h_k, l_k} \left( \sum_r (v_r^-)^2 \right) - \delta_{h_k, l_k} C(x) \geq -2 \sum_r v_r^- \delta_{h_k, l_k} v_r - \delta_{h_k, l_k} C(x),$$

which could be rewritten as

$$\sum_r \left[ v_r^- \delta_{h_k, l_k} v_r + \frac{\delta}{4(d_1+1)} \delta_{h_k, l_k} C(x) \right] \geq 0.$$

Since  $b_k \geq 0$ , this inequality implies that

$$\sum_{r,k} T_{h_r, l_r} \bar{b}_k v_r^- \delta_{h_k, l_k} v_r \geq -\frac{\delta}{4(d_1+1)} \sum_{r,k} T_{h_r, l_r} \bar{b}_k \delta_{h_k, l_k} C(x).$$

Combining this inequality with (4.12) we get the desired estimate for  $\sum_r v_r^- I_{3r}$ ,

$$(4.13) \quad \sum_r v_r^- I_{3r} \geq -\delta N(d_1, K) - N(d_1, K) M_1^2.$$

Now we consider the term  $\sum_r v_r^- I_{4r}$ .

$$\begin{aligned} \sum_r v_r^- I_{4r} &= - \sum_{r \neq \pm(d_1+1)} v_r^- \left[ (\delta_{h_r, l_r} \bar{c}) v - (T_{h_r, l_r} \bar{c}) v_r + \xi \delta_{h_r, l_r} \bar{f} \right] \\ &\quad - \sum_{r=\pm(d_1+1)} v_r^- \left[ \frac{1}{(1 \vee |l|)} (\delta_{h_r, l_r} \bar{c}) v - (T_{h_r, l_r} \bar{c}) v_r + \frac{1}{(1 \vee |l|)} \xi \delta_{h_r, l_r} \bar{f} \right]. \end{aligned}$$

We see that

$$\sum_r T_{h_r, l_r} \bar{c} (-v_r) v_r^- = \sum_r T_{h_r, l_r} \bar{c} (v_r^-)^2 \geq \lambda \sum_r (v_r^-)^2 = \lambda W.$$

Young's inequality and the definition of  $M$  then gives

$$(4.14) \quad \sum_r v_r^- I_{4r} \geq -N(K, d_1) M_1 (e^{c_0 T'} + M) + \lambda W.$$

Consider the  $\sum_r v_r^- \delta_\tau^T (\xi^{-1} v)$  term. Once more using that  $(t, x, l)$  is a maximum point of  $V$ , Lemma 4.1, and (4.1), we get

$$0 \leq -\delta_\tau^T \left( \sum_r (v_r^-)^2 \right) = -2 \sum_r v_r^- \delta_\tau^T (v_r^-) - \tau \sum_r (\delta_\tau^T (v_r^-))^2 \leq 2 \sum_r v_r^- \delta_\tau^T (v_r^-).$$

We conclude that

$$(4.15) \quad \begin{aligned} \sum_r \xi v_r^- \delta_\tau^T (\xi^{-1} v_r) &= \sum_r \xi v_r^- \left[ \xi^{-1} (t + \tau_T(t)) \delta_\tau^T v_r + v_r \delta_\tau^T \xi^{-1} \right] \\ &= e^{-c_0 \tau_T(t)} \sum_r v_r^- \delta_\tau^T v_r - W \xi \delta_\tau \xi^{-1} \\ &\geq W \frac{1 - e^{-c_0 \tau}}{\tau}. \end{aligned}$$

Using the relations (4.15), (4.14), (4.13), and (4.11), we obtain from (4.10)

$$\begin{aligned} & W\left[\frac{1 - e^{-c_0\tau}}{\tau} + \lambda\right] + \nu(E)V - \nu(E)N(d_1, K)M_1^2 - \delta N(d_1, K) - N(d_1, K)M_1^2 \\ & - N(d_1, K)M_1(e^{c_0T'} + M) + \sum_{r,k} v_r^- [\bar{a}_k \Delta_{h_k, l_k} v_r + I_{1r} + I_{2r}] \leq 0, \end{aligned}$$

i.e.,

$$\begin{aligned} & W\left[\frac{1 - e^{-c_0\tau}}{\tau} + \lambda\right] + \nu(E)V - \nu(E)N(d_1, K)M_1^2 - \delta N(d_1, K) \\ & - N(d_1, K)M_1^2 - N(d_1, K)M_1(e^{c_0T'} + M) \\ & \leq - \sum_{r,k} v_r^- \bar{a}_k \Delta_{h_k, l_k} v_r - \sum_{k, r \neq \pm(d_1+1)} v_r^- (\delta_{h_r, l_r} \bar{a}_k) \Delta_{h_k, l_k} v \\ (4.16) \quad & - \sum_{k, r = \pm(d_1+1)} \frac{1}{1 \vee |l|} v_r^- (\delta_{h_r, l_r} \bar{a}_k) \Delta_{h_k, l_k} v - \sum_{r,k} h_r v_r^- (\delta_{h_r, l_r} \bar{a}_k) \Delta_{h_k, l_k} v_r. \end{aligned}$$

Once again, using the fact that  $(t, x, l)$  is a point of maxima for  $V$ , along with the discrete product rule (Lemma 4.1) and (4.1), we have for each  $k$ ,

$$\begin{aligned} 0 & \geq \Delta_{h_k, l_k} \left( \sum_r (v_r^-)^2 \right) - \delta \Delta_{h_k, l_k} C(x) \\ & = 2 \sum_r v_r^- \Delta_{h_k, l_k} v_r^- + \sum_r [(\delta_{h_k, l_k} v_r^-)^2 + (\delta_{h_k, l_{-k}} v_r^-)^2] - \delta \Delta_{h_k, l_k} C(x) \\ & \geq -2 \sum_r v_r^- \Delta_{h_k, l_k} v_r + \sum_r [(\delta_{h_k, l_k} v_r^-)^2 + (\delta_{h_k, l_{-k}} v_r^-)^2] - \delta \Delta_{h_k, l_k} C(x). \end{aligned}$$

We rewrite this as

$$(4.17) \quad 2 \sum_r v_r^- \Delta_{h_k, l_k} v_r + \delta \Delta_{h_k, l_k} C(x) \geq \sum_r [(\delta_{h_k, l_k} v_r^-)^2 + (\delta_{h_k, l_{-k}} v_r^-)^2],$$

and conclude that

$$(4.18) \quad 2 \sum_r v_r^- \Delta_{h_k, l_k} v_r + \delta \Delta_{h_k, l_k} C(x) \geq 0.$$

Multiplying (4.17) by  $\bar{a}_k$  and summing up with respect to  $k$  we get

$$\sum_{r,k} v_r^- \bar{a}_k \Delta_{h_k, l_k} v_r + \sum_k \frac{\delta}{2} \bar{a}_k \Delta_{h_k, l_k} C(x) \geq \sum_{r,k} \bar{a}_k (\delta_{h_k, l_k} (v_r^-))^2.$$

Using this inequality, (4.16) becomes

$$\begin{aligned} & W\left[\frac{1 - e^{-c_0\tau}}{\tau} + \lambda\right] + \nu(E)V - \nu(E)N(d_1, K)M_1^2 - \delta N(d_1, K) \\ & - N(d_1, K)M_1^2 - N(d_1, K)M_1(e^{c_0T'} + M) \\ (4.19) \quad & \leq J_1 + J_2 + \frac{\delta}{4} \sum_k \bar{a}_k \Delta_{h_k, l_k} C, \end{aligned}$$

where

$$J_1 = \sum_{r,k} v_r^- |(\delta_{h_r, l_r} \bar{a}_k) \Delta_{h_k, l_k} v| - \frac{1}{4} \sum_{r,k} \bar{a}_k (\delta_{h_k, l_k} v_r^-)^2$$

$$J_2 = \sum_{r,k} v_r^- h_r |(\delta_{h_r, l_r} \bar{a}_k) \Delta_{h_k, l_k} v_r| - \frac{1}{2} \sum_{r,k} v_r^- \bar{a}_k \Delta_{h_k, l_k} v_r - \frac{1}{4} \sum_{r,k} \bar{a}_k (\delta_{h_k, l_k} v_r^-)^2.$$

Now we estimate  $J_1$ . By (4.2),

$$(4.20) \quad |\Delta_{h_k, l_k} v| \leq \sum_r |\delta_{h_k, l_k} v_r^-| + \sum_r |\delta_{h_k, l_{-k}} v_r^-|,$$

and by Lemma 4.1 and Young's inequality, we get

$$\begin{aligned} \sum_{r,k} v_r^- |(\delta_{h_r, l_r} (\bar{\sigma}_k)^2) \Delta_{h_k, l_k} v| &= \sum_{r,k} v_r^- |2\bar{\sigma}_k \delta_{h_r, l_r} \bar{\sigma}_k + h_r (\delta_{h_r, l_r} \bar{\sigma}_k)^2| \Delta_{h_k, l_k} v \\ &\leq \sum_{r,k} M_1 K |\bar{\sigma}_k \Delta_{h_k, l_k} v| + K^3 2(d_1 + 1) M_1 \sum_k h_1 |\Delta_{h_k, l_k} v| \\ &\leq N M_1 \sum_k |\bar{\sigma}_k \Delta_{h_k, l_k} v| + N M_1^2 \stackrel{(4.20)}{\leq} N \sum_{r,k} M_1 |\bar{\sigma}_k \delta_{h_k, l_k} v_r^-| + N M_1^2 \\ &\leq N \sum_{r,k} (8N M_1^2 + \frac{1}{8N} |\bar{\sigma}_k \delta_{h_k, l_k} v_r^-|^2) + N M_1^2 \leq \frac{1}{4} \sum_{r,k} \bar{a}_k (\delta_{h_k, l_k} v_r^-)^2 + N M_1^2, \end{aligned}$$

which implies that  $J_1 \leq N M_1^2$ .

The next step is to get a similar estimate on  $J_2$ . Note that

$$|a| = 2a_- + a, \quad h_r \leq K h_1, \quad h_r^2 |\Delta_{h_k, l_k} v_r| \leq M_1.$$

We get

$$\begin{aligned} &\sum_{r,k} v_r^- h_r |(\delta_{h_r, l_r} \bar{a}_k) \Delta_{h_k, l_k} v_r| \\ &\leq \sum_{r,k} v_r^- h_r |2(\delta_{h_r, l_r} \bar{\sigma}_k) \bar{\sigma}_k \Delta_{h_k, l_k} v_r + h_r (\delta_{h_r, l_r} \bar{\sigma}_k)^2 \Delta_{h_k, l_k} v_r| \\ &\leq \sum_{r,k} N_1 |v_r^- h_r \bar{\sigma}_k \Delta_{h_k, l_k} v_r| + \sum_{r,k} N_2 h_r^2 v_r^- |\Delta_{h_k, l_k} v_r| \\ &\leq \sum_{r,k} 2N_1 h_r v_r^- |\bar{\sigma}_k| (\Delta_{h_k, l_k} v_r)_- + \sum_{r,k} N_1 h_r v_r^- |\bar{\sigma}_k| |\Delta_{h_k, l_k} v_r| + N_2 M_1^2. \end{aligned}$$

In the above inequality the summation over  $r$  may be restricted to the cases where  $v_r^- \neq 0$  or  $v_r < 0$ . From (4.1) and  $h_k \Delta_{h_k, l_k} = \delta_{h_k, l_k} + \delta_{h_k, -l_k}$ , we then get

$$h_k (\Delta_{h_k, l_k} v_r)_- = h_k \max(-\Delta_{h_k, l_k} v_r, 0) \leq h_k |\Delta_{h_k, l_k} v_r^-| \leq |\delta_{h_k, l_k} v_r^-| + |\delta_{h_k, l_{-k}} v_r^-|.$$

The last two estimates give

$$\begin{aligned} &\sum_{r,k} v_r^- h_r |(\delta_{h_r, l_r} \bar{a}_k) \Delta_{h_k, l_k} v_r| \\ &\leq N_2 M_1^2 + \sum_{r,k} \left( \frac{1}{4} \bar{a}_k (\delta_{h_k, l_k} v_r^-)^2 + N_1 v_r^- h |\bar{\sigma}_k| |\Delta_{h_k, l_k} v_r| \right), \end{aligned}$$

and hence

$$J_2 \leq N_2 M_1^2 - \frac{1}{2} (\bar{a}_k - 2N_1 h \sqrt{\bar{a}_k}) v_r^- \Delta_{h_k, l_k} v_r.$$

Let

$$\mathbb{A} = \{k : (\bar{a}_k - 2N_1(K) h_1 \sqrt{\bar{a}_k}) \geq 0\}.$$

and note that if  $k \notin \mathbb{A}$ , then

$$\sqrt{\bar{a}_k} \leq 2N_1(K)h_1, \quad \bar{a}_k \leq 4N_1^2h_1^2, \quad |\bar{a}_k - 2N_1(K)h_1\sqrt{\bar{a}_k}| \leq N(K)h_1^2.$$

By (4.18) we then get

$$\begin{aligned} & -\frac{1}{2} \sum_{r,k} (\bar{a}_k - 2N_2(K)h_1\sqrt{\bar{a}_k}) v_r^- \Delta_{h_k, l_k} v_r \\ &= -\frac{1}{2} \left( \sum_{r,k \in \mathbb{A}} + \sum_{r,k \notin \mathbb{A}} \right) (\bar{a}_k - 2N_2(K)h_1\sqrt{\bar{a}_k}) v_r^- \Delta_{h_k, l_k} v_r \\ &\leq -\frac{1}{2} \sum_{r,k \in \mathbb{A}} (\bar{a}_k - 2N_2(K)h_1\sqrt{\bar{a}_k}) v_r^- \Delta_{h_k, l_k} v_r + N(K) \sum_{r,k \notin \mathbb{A}} v_r^- h_1^2 \Delta_{h_k, l_k} v_r \\ &\leq \frac{1}{2} \delta \sum_{r,k \in \mathbb{A}} (\bar{a}_k - 2N_2(K)h_1\sqrt{\bar{a}_k}) \Delta_{h_k, l_k} C(x) + N(d_1, K) M_1^2 \\ &\leq N(K, d_1) (M_1^2 + \delta), \end{aligned}$$

which gives the estimate  $J_2 \leq N(M_1^2 + \delta)$ .

The bounds on  $J_1$  and  $J_2$  along with (4.19) and the definition of  $V$  ( $V = W - \delta C$ ) give

$$\left( \lambda + \frac{1 - e^{-c_0\tau}}{\tau} \right) V(t, x, l) \leq N(\delta + (e^{c_0T'} + M_1 + M) M_1),$$

when  $t < T$ . Combining this estimate with the estimate for  $t = T$  (4.5) we see that

$$V(t, x, l) \leq \left( \lambda + \frac{1 - e^{-c_0\tau}}{\tau} \right)^{-1} N(\delta + e^{c_0T'} + M_1 + M) M_1 + N e^{2c_0T'} (\sup_{r,x} u_r(T, x))^2,$$

for every  $(t, x, l) \in \bar{\mathcal{M}}_T \times \mathbb{R}^N$  and  $\delta > 0$ . Using the definition of  $V$  and sending  $\delta \rightarrow 0$  then give for every  $t, x, l$ ,

$$\begin{aligned} & W(t, x, l) \\ (4.21) \quad & \leq \left( \lambda + \frac{1 - e^{-c_0\tau}}{\tau} \right)^{-1} N(e^{c_0T'} + M_1 + M) M_1 + N e^{2c_0T'} (\sup_{r,x} u_r(T, x))^2. \end{aligned}$$

Let  $W_{\max} = \sup_{(t,x,l) \in \bar{\mathcal{M}}_T \times \mathbb{R}^d} W(t, x, l)$ . For each  $(t, x) \in \bar{\mathcal{M}}_T$  and for each  $r$ , either  $v_r(t, x) \leq 0$  or  $-v_r(t, x) = v_{-r}(t, x + h_r l_r) \leq 0$ . In any case we have  $|v_r(t, x)| \leq \sqrt{W_{\max}}$  and hence

$$(4.22) \quad M_1 \leq \sqrt{W_{\max}} \quad \text{and} \quad \frac{1}{1 \vee |l|} |\delta_{\epsilon, \pm l} u| \leq \sqrt{W_{\max}}.$$

In view of (4.21)

$$\begin{aligned} W_{\max} &\leq \left( \lambda + \frac{1 - e^{-c_0\tau}}{\tau} \right)^{-1} N(e^{c_0T'} + \sqrt{W_{\max}} + M) \sqrt{W_{\max}} \\ &\quad + N e^{2c_0T'} (\sup_{r,x} u_r(T, x))^2. \end{aligned}$$

By this estimate, Young's inequality, and  $M \leq e^{c_0T'} |u|_0$ , we obtain

$$W_{\max} \leq \left( \lambda + \frac{1 - e^{-c_0\tau}}{\tau} \right)^{-1} N(1 + W_{\max} + |u|_0^2) + N(\sup_{r,x} u_r(T, x))^2.$$

If  $(\lambda + \frac{1-e^{-c_0\tau}}{\tau}) \geq N + 1$ , then we conclude that

$$W_{\max} \leq N' [1 + M_0^2 + (\sup_{r,x} u_r(T, x))^2].$$

Along with (4.22) this estimate proves the theorem.  $\square$

Next, following [21], we prove a continuous dependence estimate for the scheme. Let  $\hat{\sigma}_k^\alpha, \hat{b}_k^\alpha, \hat{c}^\alpha, \hat{f}^\alpha, \hat{u}_0^\alpha, \hat{\eta}^\alpha$  be functions from  $\mathcal{A} \times \mathbb{R} \times \mathbb{R}^d$  to  $\mathbb{R}$  and set  $\hat{a}_k^\alpha = \frac{1}{2} |\hat{\sigma}_k^\alpha|^2$ .

**Theorem 4.4.** *Assume (A.1), (A.3), (A.2), (3.1) and (3.2) hold. Let  $\hat{\sigma}_k^\alpha, \hat{b}_k^\alpha, \hat{c}^\alpha, \hat{f}^\alpha, \hat{\eta}^\alpha$  satisfy assumptions (A.1) – (A.3). Let  $u$  and  $\hat{u}$  be functions on  $\mathcal{M}_T$  satisfying (3.3) with coefficients  $\sigma_k^\alpha, b_k^\alpha, c^\alpha, f^\alpha, \eta^\alpha$  and  $\hat{\sigma}_k^\alpha, \hat{b}_k^\alpha, \hat{c}^\alpha, \hat{f}^\alpha, \hat{\eta}^\alpha$  respectively and  $|u(T, \cdot)|_1 + |\hat{u}(T, \cdot)|_1 \leq K$ . Define*

$$\epsilon := \sup_{\mathcal{M}_{T,A,k}} \left\{ |\hat{\sigma}_k^\alpha - \sigma_k^\alpha| + |\hat{b}_k^\alpha - b_k^\alpha| + |\hat{c}^\alpha - c^\alpha| + |\hat{f}^\alpha - f^\alpha| + |\hat{\eta}^\alpha - \eta^\alpha| \right\}.$$

Then if there exists a  $c_0 \geq 0$  satisfying (4.4), there exists a constant  $N$  depending only on  $K, d_1, \lambda, c_0, T, \nu(E)$  such that

$$(4.23) \quad |u - \hat{u}| \leq NI\epsilon \quad \text{on } \bar{\mathcal{M}}_T,$$

where

$$I := 1 + \max_k |\delta_{h_1, l_k} u(T, \cdot)|_0 + \max_k |\delta_{h_1, l_k} \hat{u}(T, \cdot)|_0 + \epsilon^{-1} |u(T, \cdot) - \hat{u}(T, \cdot)|_0.$$

*Proof.* First we show that it is sufficient to prove the result assuming  $\epsilon \leq h_1$ . For each  $\theta \in [0, 1]$ , let  $u^\theta$  be the (unique) solution of

$$\begin{aligned} \delta_\tau^T u + \sup_{\alpha \in \mathcal{A}} [a_k^{\theta\alpha} \Delta_{h_1, l_k} u + b_k^{\theta\alpha} \delta_{h_k, l_k} u - c^{\theta\alpha} u + f^{\theta\alpha} \\ + \sum_p k_p (u(t, x + \eta^{\theta\alpha}(x, p)) - u(t, x))] = 0 \quad \text{in } \mathcal{M}_T, \end{aligned}$$

with  $u^\theta(T, x) = (1 - \theta)u(T, x) + \theta\hat{u}(T, x)$  and where

$$[a_k^{\theta\alpha}, b_k^{\theta\alpha}, c^{\theta\alpha}, f^{\theta\alpha}, \eta^{\theta\alpha}] = (1 - \theta)[\sigma_k^\alpha, b_k^\alpha, c^\alpha, f^\alpha, \eta^\alpha] + \theta[\hat{\sigma}_k^\alpha, \hat{b}_k^\alpha, \hat{c}^\alpha, \hat{f}^\alpha, \hat{\eta}^\alpha].$$

By uniqueness,  $u^0 = u$  and  $u^1 = \hat{u}$ . Also note that for any  $\theta_1, \theta_2 \in [0, 1], \alpha, k$ ,

$$\begin{aligned} |\sigma_k^{\theta_1\alpha} - \sigma_k^{\theta_2\alpha}|_0 + |b_k^{\theta_1\alpha} - b_k^{\theta_2\alpha}|_0 + |c^{\theta_1\alpha} - c^{\theta_2\alpha}|_0 + |f^{\theta_1\alpha} - f^{\theta_2\alpha}|_0 + |\eta^{\theta_1\alpha} - \eta^{\theta_2\alpha}|_0 \\ \leq |\theta_1 - \theta_2|\epsilon. \end{aligned}$$

Therefore if we assume the result holds for  $\epsilon \leq h_1$ , then for any  $\epsilon$  satisfying  $|\theta_1 - \theta_2|\epsilon \leq h_1$ , we have

$$(4.24) \quad |u^{\theta_1} - u^{\theta_2}| \leq N_1 |\theta_1 - \theta_2| \epsilon I(\theta_1, \theta_2)$$

where

$$\begin{aligned} I(\theta_1, \theta_2) = 1 + \max_k |\delta_{h_1, l_k} u^{\theta_1}(T, \cdot)|_0 + \max_k |\delta_{h_1, l_k} u^{\theta_2}(T, \cdot)|_0 \\ + \epsilon^{-1} |\theta_1 - \theta_2|^{-1} |u^{\theta_1}(T, \cdot) - u^{\theta_2}(T, \cdot)|_0. \end{aligned}$$

Clearly  $I(\theta_1, \theta_2) \leq 4I$ , so by dividing the interval  $[0, 1]$  into sufficient number of sub-intervals  $\theta_1, \dots, \theta_n$ , we can conclude the theorem (with  $4N$  instead of  $N$ ), by writing

$$u - \hat{u} = \sum_{i=1}^n (u^{\theta_i} - u^{\theta_{i-1}}),$$

and using (4.24) to estimate each  $u^{\theta_i} - u^{\theta_{i-1}}$ . Henceforth we assume that  $\epsilon \leq h_1$ .

We will now show that the continuous dependence estimate (4.23) is a consequence of the Lipschitz estimate Theorem (4.3). To this end, we consider  $\mathbb{R}^d$  as subspace of  $\mathbb{R}^{d+1}$  and write,

$$\begin{aligned} \mathbb{R}^{d+1} &= \{(x', x^{d+1}) : x' \in \mathbb{R}^d, x^{d+1} \in \mathbb{R}\}, \\ \bar{Q}_T(d+1) &= [0, T] \times \mathbb{R}^{d+1}, \quad \bar{\mathcal{M}}_T(d+1) = \{j\tau \wedge T : j = 0, 1, 2, \dots\} \times \mathbb{R}^{d+1}, \\ Q_T(d+1) &= [0, T] \times \mathbb{R}^{d+1}, \quad \text{and} \quad \mathcal{M}_T(d+1) = Q_T(d+1) \cap \bar{\mathcal{M}}_T(d+1). \end{aligned}$$

Let  $\rho \in C^1(\mathbb{R})$  be a bounded function on  $\mathbb{R}$  such that

$$\rho(-1) = 1, \quad \rho(0) = 0, \quad \rho'(p) = \rho'(q) = 0 \text{ if } p \leq -1, \quad q \geq 0.$$

Now define

$$\tilde{\sigma}_k^\alpha(t, x', x_{d+1}) := \tilde{\sigma}_k^\alpha(t, x') \rho\left(\frac{x^{d+1}}{\epsilon}\right) + \sigma_k^\alpha(t, x') \left(1 - \rho\left(\frac{x^{d+1}}{\epsilon}\right)\right),$$

and in a similar way,  $\tilde{b}_k^\alpha, \tilde{c}_k^\alpha, \tilde{f}_k^\alpha, \tilde{u}(T, \cdot)$ . Define  $\tilde{\eta}^\alpha : \mathbb{R}^{d+1} \times \mathbb{R}^M \mapsto \mathbb{R}^{d+1}$  as

$$\tilde{\eta}^\alpha(x', x_{d+1}; z) = (\hat{\eta}^\alpha(x', z), 0) \rho\left(\frac{x^{d+1}}{\epsilon}\right) + (\eta^\alpha(x', z), 0) \left(1 - \rho\left(\frac{x^{d+1}}{\epsilon}\right)\right).$$

We would like to show that,  $\tilde{\sigma}_k^\alpha, \tilde{b}_k^\alpha, \tilde{c}_k^\alpha, \tilde{f}_k^\alpha, \tilde{\eta}^\alpha$  and  $\tilde{u}(T, \cdot)$  all satisfy the assumptions (A.1) and (A.3) in  $\bar{Q}_T(d+1)$ . All other properties apart from Lipschitz continuity in  $(d+1)$ -th direction are straight forward. For  $\tilde{\sigma}_k^\alpha$  along  $(d+1)$ -direction we have

$$\left| \frac{\partial \tilde{\sigma}_k^\alpha}{\partial x^{d+1}} \right| = \frac{1}{\epsilon} |\tilde{\sigma}_k^\alpha(t, x') - \sigma_k^\alpha(t, x')| \left| \rho'\left(\frac{x^{d+1}}{\epsilon}\right) \right| \leq \left| \rho'\left(\frac{x^{d+1}}{\epsilon}\right) \right| \leq K.$$

A similar conclusion holds for the other functions.

Therefore by Lemma 3.1 there exists a function  $\tilde{u}_{\tau, h}$ , defined on  $\bar{\mathcal{M}}_T(d+1)$ , which solves (3.3) with the new family of coefficients  $\tilde{\sigma}_k^\alpha, \tilde{b}_k^\alpha, \tilde{c}_k^\alpha, \tilde{f}_k^\alpha, \tilde{\eta}^\alpha$  and terminal data  $\tilde{u}(T, \cdot)$ . Furthermore, by uniqueness, we must have

$$\tilde{u}_{\tau, h}(t, x', -\epsilon) = \hat{u}(t, x'), \quad \tilde{u}_{\tau, h}(t, x', 0) = \tilde{u}_{\tau, h}(t, x', \epsilon) = u(t, x').$$

Now we choose  $l = (0, 0, \dots, 1)$ , the unit vector along the  $(d+1)$ -th direction. For this  $l$ , by Theorem 4.3 (since  $\epsilon \leq h_1$ ) we conclude that there exists a constant  $N$ , depending only on  $d, d_1, K, c_0, \lambda, T$ , and  $\nu$ , such that

$$\left| \frac{\tilde{u}_{\tau, h}(t, x', -\epsilon) - \tilde{u}_{\tau, h}(t, x', 0)}{\epsilon} \right| \leq N \sup_{k, x, l} \left[ 1 + |\delta_{h_1, l_k} \tilde{u}(T, x)| + \frac{1}{1 \vee |l|} |\delta_{\epsilon, \pm l} \tilde{u}(T, x)| \right],$$

which gives  $|\hat{u} - u| \leq NI\epsilon$ .  $\square$

Next we establish Lipschitz continuity property of  $v_{\tau, h}$  in the  $x$ -variable. To do this, let  $S \subset B_1 = \{x \in \mathbb{R}^d : |x| < 1\}$  be nonempty,  $\epsilon \in \mathbb{R}$ , and  $v_{\tau, h}^{\epsilon, S}$  be the unique solution of the equation

$$\begin{aligned} \delta_\tau^T u + \sup_{(\alpha, y) \in \mathcal{A} \times S} & \left[ L_{h_1}^\alpha(t, x + \epsilon y) u(t, x) + f^\alpha(t, x + \epsilon y) \right. \\ & \left. + \sum_p k_p (u(t, x + \eta^\alpha(x + \epsilon y, p)) - u(t, x)) \right] = 0 \quad \text{in } Q_T, \end{aligned}$$

with the terminal data

$$(4.25) \quad u(T, x) = \sup_{y \in S} u_0(x + \epsilon y).$$

Note that if  $S$  is a singleton  $\{y\}$ , then by uniqueness  $v_{\tau,h}^{\epsilon,S}(t,x) = v_{\tau,h}(t,x + \epsilon y)$ .

**Lemma 4.5.** *Assume (A.1), (A.3), (A.2), and (3.1),(3.2) hold. There is a constant  $N$  depending only on  $K, d_1, \nu(E), T$ , so that if the condition (4.4) is satisfied for a constant  $c_0 \geq 0$ , then for every  $\epsilon \in \mathbb{R}$ ,*

$$(4.26) \quad |v_{\tau,h}^{\epsilon,S} - v_{\tau,h}| \leq N_1 |\epsilon| \quad \text{on } \bar{Q}_T,$$

where  $N_1$  only depend on  $K, d_1, \nu(E), T, \lambda, c_0$ . Furthermore, by choosing  $S := \{\frac{y-x}{|y-x|}\}$ ,  $\epsilon = |y-x|$ , we have

$$(4.27) \quad |v_{\tau,h}(t,x) - v_{\tau,h}(t,y)| \leq N_1 |y-x| \quad \text{for all } (t,y), (t,x) \in \bar{Q}_T.$$

*Proof.* It is sufficient to prove (4.26) on  $\bar{M}_T$ . We use of Theorem 4.4, where we choose  $A \times S, (\sigma, b, c, f, \eta)$  and  $(\sigma, b, c, f, \eta)(x + \epsilon y, t/z)$  in places of  $A, (\sigma, b, c, f, \eta)$  and  $\hat{\sigma}, \hat{b}, \hat{c}, \hat{f}, \hat{\eta}$ , respectively. The contribution from the difference of the terminal data can be bounded by  $N\epsilon$ .  $\square$

A step in the direction of establishing Theorem 3.4 is to prove

**Lemma 4.6.** *Assume (A.1), (A.2), (A.3) and (3.1),(3.2) hold. Let  $h_1, h_2, \tau \leq K$ . Let  $(s_0, x_0) \in \bar{M}_T$  and set*

$$L := \sup_{x \in \mathbb{R}^N, x \neq x_0} \frac{|v_{\tau,h}(s_0, x) - v_{\tau,h}(s_0, x_0)|}{|x - x_0|}.$$

Then for all  $(t_0, x_0) \in \mathcal{M}_T$  satisfying  $s_0 - 1 \leq t_0 \leq s_0$  and  $\frac{1}{\tau}T \in \mathbb{N}$ , we have

$$|v_{\tau,h}(s_0, x_0) - v_{\tau,h}(t_0, x_0)| \leq N(L + 1)|s_0 - t_0|^{\frac{1}{2}},$$

where  $N$  depends only on  $K, d_1$  and  $\nu(E)$ .

*Proof.* Without loss of generality we may restrict ourselves to the case  $0 < s_0 < 1$  and  $t_0 = 0$ . This claim follows by shifting the origin and observing that there holds  $|v_{\tau,h}(s_0, x_0) - v_{\tau,h}(t_0, x_0)| \leq 2|v_{\tau,h}|_0 |s_0 - t_0|^{\frac{1}{2}}$  whenever  $|t_0 - s_0| > 1$ .

Fix a constant  $\gamma > 0$ . We are going to work with  $\bar{M}_{s_0}$ . On  $\bar{M}_{s_0}$  we define

$$\psi = \gamma L \left[ \zeta + \kappa(s_0 - t) \right] + K(s_0 - t) + \gamma^{-1}L + v_{\tau,h}(s_0, x_0),$$

where

$$\xi(t) = e^{s_0 - t}, \quad \beta(x) = |x - x_0|^2, \quad \zeta = \xi\eta,$$

and the constant  $\kappa$  will be chosen later. We will show that if  $\kappa$  is big enough, then  $\psi$  is a supersolution of (3.3).

On  $M_{s_0}$  we have,  $\delta_\tau^{s_0} \xi = -\theta \xi$  where,  $\theta := \tau^{-1}(1 - e^{-\tau}) \geq K^{-1}(1 - e^{-K})$ . Furthermore, using (A.1) – (A.3) we get

$$\begin{aligned} & \mathcal{L}_{h_1}^\alpha \beta(t, x) + \sum_p k_p [\beta(t, x + \eta^\alpha(x, p)) - \beta(t, x)] \\ &= 2a_k^\alpha(t, x) |l_k|^2 + b_k^\alpha(t, x) (l_k, 2(x - x_0) + h_1 l_k) - c^\alpha \beta(t, x) \\ & \quad + \sum_p k_p \langle \eta^\alpha(x, p); 2(x - x_0) + \eta^\alpha(x, p) \rangle \\ & \leq N_1(d_1, K, \nu(E))(1 + |x - x_0|), \end{aligned}$$

and hence

$$\begin{aligned} & \delta_\tau^{s_0} \zeta(t, x) + \mathcal{L}_{h_1}^\alpha \zeta(t, x) + \sum_p k_p [\zeta(t, x + \eta^\alpha(x, p)) - \zeta(t, x)] \\ & \leq N(d_1, K, \nu(E))(1 + |x - x_0|) - \theta(\tau)|x - x_0|^2. \end{aligned}$$

Applying the same operator on  $\psi$  and using the above estimates, we have

$$\begin{aligned} & \delta_\tau^{s_0} \psi + \mathcal{L}_{h_1}^\alpha \psi + f^\alpha + \sum_p k_p [\psi(t, x + \eta^\alpha(x, p)) - \psi(t, x)] \\ & \leq \gamma L [N_2(1 + |x - x_0|) - \theta|x - x_0|^2 - \kappa] + f^\alpha - K. \end{aligned}$$

Since  $|f^\alpha| \leq K$  and by suitably applying Young's inequality:  $2ab \leq ra^2 + \frac{b^2}{r}$ , it is clear that there exist  $\kappa$  depending only on  $N_2$  such that the right hand side of the last inequality is negative. So, for this choice of  $\kappa$  we have

$$\delta_\tau^{s_0} \psi + \sup_{\alpha \in \mathfrak{a}} [\mathcal{L}_{h_1}^\alpha \psi + f^\alpha + \sum_p k_p [\psi(t, x + \eta^\alpha(x, p)) - \psi(t, x)]] \leq 0$$

and

$$\begin{aligned} \psi(s_0, x) &= L(\gamma|x - x_0|^2 + \gamma^{-1}) + v_{\tau, h}(s_0, x_0) \\ &\geq L|x - x_0| + v_{\tau, h}(s_0, x_0) \geq v_{\tau, h}(s_0, x). \end{aligned}$$

We now apply Lemma 3.2 on  $M_{s_0}$  and conclude that

$$v_{\tau, h}(t, x_0) \leq \psi(t, x_0) = \gamma L \kappa (s_0 - t) + \gamma^{-1} L + K(s_0 - t) + v_{\tau, h}(s_0, x_0).$$

Minimizing with respect to the  $\gamma > 0$  and using the fact  $(s_0 - t) \leq 1$ , we conclude

$$\begin{aligned} v_{\tau, h}(t, x_0) - v_{\tau, h}(s_0, x_0) &\leq 2L\kappa^{\frac{1}{2}}|s_0 - t|^{\frac{1}{2}} + Ks_0^{\frac{1}{2}}|s_0 - t|^{\frac{1}{2}} \\ &\leq N(d_1, T, K, \nu(E))(L + 1)|s_0 - t|^{\frac{1}{2}}. \end{aligned}$$

The estimate for the other side is obtained similarly.  $\square$

We close this section by giving the proof of Theorem 3.4.

**Lemma 4.7.** *Assume (A.1), (A.2), (A.3), (3.1), (3.2),  $h_1, h_2, \tau \leq K$ , and*

$$L := \sup_{(t, x), (t, y) \in \bar{Q}_\tau, x \neq y} \frac{|v_{\tau, h}(t, x) - v_{\tau, h}(t, y)|}{|x - y|}.$$

Then we have

$$|v_{\tau, h}(s, x) - v_{\tau, h}(t, x)| \leq N(1 + L)(|s - t|^{\frac{1}{2}} + \tau^{\frac{1}{2}}),$$

where  $N$  depends only on  $K, d_1$  and  $\nu(E)$ .

*Proof.* If  $|t - s| \geq 1$ , then  $|v_{\tau, h}(t, x) - v_{\tau, h}(s, x)| \leq 2|v_{\tau, h}|_0 |t - s|^{\frac{1}{2}}$ . We may therefore assume  $|t - s| \leq 1$ .

Assume (without loss of generality) that  $s - t = n\tau + \gamma$  where  $\gamma \in [0, \tau)$  and  $n$  is a natural number. If  $\gamma = 0$ , then we apply Lemma 4.6 on  $(t, 0) + \mathcal{M}_{n\tau}$  and conclude

$$(4.28) \quad |v_{\tau, h}(t, x) - v_{\tau, h}(t + n\tau, x)| \leq N|n\tau|^{\frac{1}{2}}.$$

Now, for other case when  $0 < \gamma < \tau$  we have,

$$\begin{aligned} |v_{\tau, h}(t, x) - v_{\tau, h}(s, x)| &\leq |v_{\tau, h}(t, x) - v_{\tau, h}(t + n\tau, x)| + |v_{\tau, h}(s - \gamma, x) - v_{\tau, h}(s, x)| \\ &\leq N|n\tau|^{\frac{1}{2}} + |v_{\tau, h}(s - \gamma, x) - v_{\tau, h}(s, x)|. \end{aligned}$$

Therefore, we have to estimate  $|v_{\tau,h}(s-\gamma, x) - v_{\tau,h}(s, x)|$  for  $\gamma \in (0, \tau)$  and  $s-\gamma \geq 0$ . Define the following functions on  $(s, 0) + \bar{\mathcal{M}}_{T-s}$ :

$$[\hat{\sigma}_k^\alpha, \hat{b}_k^\alpha, \hat{c}^\alpha, \hat{f}^\alpha](r, y) = [\sigma_k^\alpha, b_k^\alpha, c^\alpha, f^\alpha](r - \gamma, y)$$

and

$$u = v_{\tau,h}; \hat{u}(r, y) = v_{\tau,h}(r - \gamma, y),$$

for all  $(r, y) \in (s, 0) + \bar{\mathcal{M}}_{T-s}$ . Then, on  $(s, 0) + \bar{\mathcal{M}}_{T-s}$ ,  $\hat{u}$  satisfies (3.3) constructed from  $\hat{\sigma}_k^\alpha, \hat{b}_k^\alpha, \hat{c}^\alpha, \hat{f}^\alpha$  and unchanged jump amplitudes  $\eta^\alpha$ . Noticing the fact that the parameter  $\epsilon$  in Theorem 4.4, after using (A.3), is less than  $N\gamma^{\frac{1}{2}}$  and also using the  $x$ -Lipschitz continuity of  $v_{\tau,h}$  we have, after applying Theorem 4.4 on  $(s, 0) + \bar{\mathcal{M}}_{T-s}$ ,

$$(4.29) \quad \begin{aligned} |v_{\tau,h}(s, x) - v_{\tau,h}(s - \gamma, y)| &= |u(s, x) - \hat{u}(s, y)| \\ &\leq N\gamma^{\frac{1}{2}} + \sup_{y \in \mathbb{R}^d} |u(T, y) - \hat{u}(T, y)| \\ &= N\gamma^{\frac{1}{2}} + |v_{\tau,h}(T, x) - v_{\tau,h}(T - \gamma, y)|. \end{aligned}$$

Lastly, we are left with estimating  $|v_{\tau,h}(T, x) - v_{\tau,h}(T - \gamma, y)|$ . To this end, consider the grid  $\mathcal{M}_\tau$ . With a slight abuse of the notation, we define  $\hat{u}(r, x)$  on  $\bar{\mathcal{M}}_\tau$  by

$$\hat{u}(0, x) = v_{\tau,h}(T - \gamma, x); \hat{u}(\tau, x) = v_{\tau,h}(T, x).$$

Then  $\hat{u}$  solves (3.3) on  $\mathcal{M}_\tau$ , with an obvious shift (by a quantity  $t-\gamma$  in the backward direction) in the time variable of the coefficients, and therefore by Lemma 4.6 we must have

$$|\hat{u}(\tau, x) - \hat{u}(0, x)| \leq N\tau^{\frac{1}{2}},$$

i.e.,

$$(4.30) \quad |v_{\tau,h}(T, x) - v_{\tau,h}(T - \gamma, x)| \leq N\tau^{\frac{1}{2}}.$$

Finally, we conclude by combining the estimates (4.28), (4.29), and (4.30).  $\square$

*Proof of Theorem 3.4.* Estimates (3.8), (3.9) follow from Theorem 4.4, and estimate (3.10) from Lemma 4.7 if  $\tau$  is small enough (then  $L < \infty$  by Theorem 4.3).  $\square$

## 5. SINGULAR LÉVY MEASURES AND OPTIMAL ERROR BOUNDS IN A SPECIAL CASE

In this section we address the case of unbounded (singular) Lévy measures. Specifically, in a special case, we introduce a modified difference-quadrature scheme for which we obtain an optimal (under our assumptions) convergence rate.

In general the Lévy measure need not be bounded and/or compactly supported, but it always satisfies the condition

$$(5.1) \quad \int_{|z| \leq 1} |z|^2 \nu(dz) + \int_{|z| > 1} e^{K|z|} \nu(dz) < \infty,$$

for some constant  $K \geq 0$ . Under this condition, the jump amplitude  $\eta^\alpha$  must satisfy

$$(5.2) \quad |\eta^\alpha(x, z)| + \sup_{|h| > 0} \frac{1}{h} |\eta^\alpha(x + h, z) - \eta^\alpha(x, z)| \leq N(|z| 1_{|z| \leq 1} + e^{K|z|} 1_{|z| > 1}).$$

Conditions (5.1) and (5.2) along with (A.1) and (A.3) ensure that the underlying stochastic control problem is well-defined. Moreover, the initial value problem (1.1) and (1.2), with  $\mathcal{I}$  defined in (1.3), possesses a unique Hölder continuous viscosity

solution. We refer to [16] for the proof of this result and for the precise definition of viscosity solutions in this setting.

To solve such problems numerically the first step is often (see, e.g., [12, 18]) to approximate the Lévy measure  $\nu$  by a finite and compactly supported measure of the form  $\nu_{r,R}(z) = \mathbf{1}_{r < |z| < R} \nu(dz)$  (occasionally one also adds a diffusion term to the equation to account for the small jumps  $|z| \leq r$ ), and then to discretize the corresponding Bellman equation by a finite difference-quadrature scheme like (3.3) of Section 3. The truncation error related to  $r, R$  can be estimated following the arguments of [18], while for a fixed truncation level, i.e., for a fixed choice of  $r$  and  $R$ , the error coming from the numerical scheme is given by Theorem 3.5.

By choosing  $r, R$  optimally in terms of  $\tau$  and  $h$ , it should, at least in principle, be possible to derive a convergence rate for this scheme. However, this is not straightforward since the error bound of Theorem 3.5 only holds if either  $\lambda$  is sufficiently large or  $\tau$  is sufficiently small when  $r$  is small (and  $\nu_{r,R}(E)$  is large), see the proofs of Theorems 3.5 and 4.3. This difficulty can most likely be overcome, for example, by an iteration argument of the type used to prove the Hölder estimates in [4]. Perhaps more important, such a convergence rate cannot be optimal because the Lipschitz estimate in Theorem 4.3 is not optimal when the Lévy measure is singular (it “deteriorates” as  $r \rightarrow 0$ ).

In the remaining part of this section we will present a different approach, which in the end will produce a better convergence rate. This approach is based on a direct discretization of the integral term (1.3) (no truncation) and obtaining an optimal Lipschitz estimates for the corresponding numerical scheme. The main idea is to use a more refined Bernstein argument than the one used to prove Theorem 4.3, one that mimics the Bernstein argument for the continuous integro-PDE (with an singular Lévy measure). In implementing this idea we shall restrict ourselves to the case where the jumps do not depend on  $x$ :

$$\eta^\alpha(x, z) = \eta^\alpha(z) \quad (\text{i.e., } \eta^\alpha \text{ does not depend on } x).$$

This assumption (along with previous ones) are sufficient to imply a Lipschitz estimate for our modified numerical scheme that applies when the Lévy measure is singular (see below). To obtain a convergence rate for our scheme, we will impose an additional technical condition saying that for a fixed  $\alpha$  it is possible to jump only in one direction, i.e.,

$$(5.3) \quad \eta^\alpha(x, z) = \xi^\alpha \eta^\alpha(z),$$

for some direction  $\xi^\alpha \in \mathbb{R}^N$ ,  $|\xi^\alpha| = 1$ , and a scalar function  $\eta^\alpha(z)$  satisfying (5.2). This restrictive assumption is used to ensure that the integrand in the unbounded Lévy case is Lipschitz continuous in  $z$  (details are given below).

Let us now turn to the precise definition our scheme. To this end, we introduce the finite measures

$$\nu^\alpha(dz) := \mathbf{1}_{|z| \leq 1} |\eta^\alpha(z)|^2 \nu(dz) + \mathbf{1}_{|z| \geq 1} \nu(dz).$$

These positive measures are nonsingular at the origin. Let  $I_{h_2}^\alpha$  be a quadrature rule satisfying

$$(5.4) \quad I_{h_2}^\alpha f = \sum_{p \in h_2 \mathbb{Z}^N \setminus \{0\}} k_p^\alpha f(p), \quad \text{where } k_p^\alpha \geq 0,$$

and

$$(5.5) \quad |I_{h_2}^\alpha f - \int_E f(z) \nu^\alpha(dz)| \leq L_f \max_\alpha \nu^\alpha(E) h_2,$$

for bounded Lipschitz continuous functions  $f(z)$  with Lipschitz constant  $L_f$ .

Conditions 5.4 and 5.5 are satisfied by Riemann sum approximations and Newton-Cotes quadratures of order less than 9 like the compound trapezoidal and Simpson rules. Note that by (5.5),  $I_{h_2} 1 = \sum_p k_p^\alpha = \nu^\alpha(E)$  and that  $\max_\alpha \nu^\alpha(E) < \infty$  by (5.1) and (5.2).

We discretize the non-local term in (1.1) in the following way:

$$\mathcal{I}_{h_2}^\alpha u(t, x) := I_{h_2}^\alpha \left[ \frac{u(t, x + \xi^\alpha \eta^\alpha(z)) - u(t, x) + 1_{|z| \leq 1} \frac{1}{h_1} (u(t, x - h_1 \xi^\alpha \eta^\alpha(z)) - u(t, x))}{1_{|z| \leq 1} |\eta^\alpha(z)|^2 + 1_{|z| > 1}} 1_{\{\eta^\alpha \neq 0\}} \right].$$

Our new difference-quadrature scheme for (1.1) now reads

$$(5.6) \quad \delta_\tau^T u(t, x) + \sup_{\alpha \in \mathcal{A}} \left[ \mathcal{L}_{h_1}^\alpha(t, x) u + f^\alpha(t, x) + \mathcal{I}_{h_2}^\alpha u \right] = 0.$$

Existence, uniqueness, and comparison results for (5.6) follow along the same lines as before. Furthermore, we claim that the Lipschitz estimate of Theorem 3.4 is satisfied for (5.6) with the modification that the constant  $\tau_0$  along with the different constants  $N$  appearing in (3.8), (3.9), (3.10) are independent of  $\nu(E)$ . To verify this claim, it is enough to show that Theorem 4.3 can be proved with  $N$  in (4.4) chosen independent of  $\nu(E)$ .

The “ $\nu(E)$  - relevant” term in the proof of Theorem 4.3 is  $v_r^- I_{5r}$  which in the present context equals  $v_r^- \mathcal{I}_{h_2}^\alpha v_r^-$ . With a slight abuse of the notation we define

$$I_h w(x) := w(x + \eta(z)) - w(x) + 1_{|z| \leq 1} \frac{1}{h_1} [w(x - h_1 \eta^\alpha(z)) - w(x)],$$

for bounded functions  $w(x)$ . If, at a point  $x$ ,  $w(x) \leq 0$ , then

$$(5.7) \quad \begin{aligned} I_h w(x) &:= w(x + \eta(z)) - w(x) + 1_{|z| \leq 1} \frac{1}{h_1} [w(x - h_1 \eta^\alpha(z)) - w(x)] \\ &\geq -w^-(x + \eta(z)) + w^- - 1_{|z| \leq 1} \frac{1}{h_1} [w^-(x - h_1 \eta^\alpha(z)) - w^-(x)] \\ &= -I_h(w^-)(x). \end{aligned}$$

Since  $(t, x)$  is a maximum point of  $V$  (remember that we are following the proof of Theorem 4.3), we have

$$\begin{aligned} 0 &\geq I_h \sum_r (v_r^-)^2 - \delta I_h C(x) \\ &= 2v_r^- I_h v_r^- + \text{“positive quantity”} - \delta I_h C(x), \end{aligned}$$

and, since the sum over  $r$  could be restricted to those  $r$  for which  $v_r \leq 0$ , it follows from (5.7) that

$$0 \geq -2v_r^- I_h v_r^- - \delta I_h C(x).$$

Hence, by the obvious relation between  $I_{h_2}$  and  $\mathcal{I}_{h_2}^\alpha$ ,

$$v_r^- I_{5r} = v_r^- \mathcal{I}_{h_2}^\alpha v_r^- \geq -\frac{\delta}{2} \mathcal{I}_{h_2}^\alpha C(x) \geq -\frac{\delta}{2} N,$$

where  $N$  does not depend on  $\nu(E)$ . This proves the claim.

For any smooth function  $w$ , let us define the function

$$G(z) = \frac{w(x + \xi^\alpha \eta^\alpha(z)) - w(x) - \xi^\alpha \eta^\alpha(z) D_x w}{|\eta^\alpha(z)|^2},$$

and observe that

$$|D_z G| \leq 2|(\eta^\alpha)'|_0 |D_x^3 w|_0,$$

where  $(\eta^\alpha)'$  is the  $z$ -derivative of  $\eta^\alpha$ . Condition (5.3) is used to ensure the validity of this estimate; For more general forms of  $\eta^\alpha(z)$  it may not be true. As a consequence of this bound and a split of the domain of integration into the two regions  $\{|z| \leq 1\}$  and  $\{|z| > 1\}$ , we obtain the error bound

$$(5.8) \quad |\mathcal{I}^\alpha w - \mathcal{I}_h^\alpha w| \leq N((|D_x^3 w|_0 + |D_x w|_0)h_2 + |D_x^2 w|_0 h_1).$$

Denote by  $v_{\tau,h}$  the unique solution of (5.6), (1.2) and by  $v$  the unique viscosity solution of (1.1), (1.2). In view of (5.8) and the discussion above, if we repeat the proof of Theorem 3.5 we will eventually find that

$$|v - v_{\tau,h}| \leq N \min\left(\epsilon + h_2\left(\frac{1}{\epsilon} + \frac{1}{\epsilon^2}\right) + \frac{h_1}{\epsilon} + \frac{\tau + h_1^2}{\epsilon^3}\right) \leq N(\tau^{\frac{1}{4}} + h_1^{\frac{1}{2}} + h_2^{\frac{1}{3}}),$$

for sufficiently small  $\tau, h_1, h_2$ .

Summarizing, we have proved

**Theorem 5.1.** *Assume that (A.1), (A.3), (5.1), (5.2), (5.3), (5.4), (5.5) hold. Let  $v$  and  $v_{\tau,h}$  be the solutions of (1.1), (1.2) and (5.6), (1.2), respectively. Then there exists a constant  $N$ , depending only on  $d, d_1$ , and  $K$ , such that*

$$|v - v_{\tau,h}| \leq N(\tau^{\frac{1}{4}} + h_1^{\frac{1}{2}} + h_2^{\frac{1}{3}}).$$

**Remark 5.1.** Theorem 5.1 appears to be the first result on convergence rates for numerical schemes of Bellman equations with singular Lévy measures. Note that the convergence rate in Theorem 5.1 does not depend on the strength of the singularity at  $z = 0$  of the Lévy measure. Furthermore, the result is most likely optimal under the current conditions. However, if we have further information about  $\nu$ , e.g., if  $\nu(dz) \leq N|z|^{-\gamma-M} dz$  in a neighborhood of  $z = 0$  for some  $\gamma \in (0, 2)$ , then the rate  $1/3$  can be improved. We leave this to the interested reader.

## REFERENCES

- [1] O. Alvarez and A. Tourin. Viscosity solutions of nonlinear integro-differential equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 13(3):293–317, 1996.
- [2] A. L. Amadori. Nonlinear integro-differential evolution problems arising in option pricing: a viscosity solutions approach. *Differential Integral Equations*, 16(7):787–811, 2003.
- [3] G. Barles, R. Buckdahn, and E. Pardoux. Backward stochastic differential equations and integral-partial differential equations. *Stochastics Stochastics Rep.*, 60(1-2):57–83, 1997.
- [4] G. Barles and E. R. Jakobsen. On the convergence rate of approximation schemes for Hamilton-Jacobi-Bellman equations. *M2AN Math. Model. Numer. Anal.*, 36(1):33–54, 2002.
- [5] G. Barles and E. R. Jakobsen. Error bounds for monotone approximation schemes for Hamilton-Jacobi-Bellman equations. *SIAM J. Numer. Anal.*, 43(2):540–558 (electronic), 2005.
- [6] G. Barles and E. R. Jakobsen. Error bounds for monotone approximation schemes for parabolic Hamilton-Jacobi-Bellman equations. *Math. Comp.*, 76(240):1861–1893, 2007.
- [7] G. Barles and P. E. Souganidis. Convergence of approximation schemes for fully nonlinear second order equations. *Asymptotic Anal.*, 4(3):271–283, 1991.

- [8] F. E. Benth, K. H. Karlsen, and K. Reikvam. Optimal portfolio management rules in a non-Gaussian market with durability and intertemporal substitution. *Finance Stoch.*, 5(4):447–467, 2001.
- [9] F. E. Benth, K. H. Karlsen, and K. Reikvam. Optimal portfolio selection with consumption and nonlinear integrodifferential equations with gradient constraint: a viscosity solution approach. *Finance and Stochastic*, 5:275–303, 2001.
- [10] F. E. Benth, K. H. Karlsen, and K. Reikvam. Portfolio optimization in a Lévy market with intertemporal substitution and transaction costs. *Stoch. Stoch. Rep.*, 74(3-4):517–569, 2002.
- [11] I. H. Biswas, E. R. Jakobsen, and K. H. Karlsen. Error estimates for finite difference-quadrature schemes for a class of nonlocal Bellman equations with variable diffusion. *Contemp. Math.*, volume 429, Amer. Math. Soc., pages 19-31, 2007.
- [12] R. Cont and P. Tankov. *Financial modeling with jump processes*. Chapman & Hall/CRC, Boca Raton, FL, 2004.
- [13] M. G. Crandall, H. Ishii, and P.-L. Lions. User’s guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc. (N.S.)*, 27(1):1–67, 1992.
- [14] W. H. Fleming and H. M. Soner. *Controlled Markov processes and viscosity solutions*. Springer-Verlag, New York, 1993.
- [15] E. R. Jakobsen. On the rate of convergence of approximation schemes for Bellman equations associated with optimal stopping time problems. *Math. Models Methods Appl. Sci.*, 13(5):613–644, 2003.
- [16] E. R. Jakobsen and K. H. Karlsen. Continuous dependence estimates for viscosity solutions of integro-PDEs. *J. Differential Equations*, 212(2):278–318, 2005.
- [17] E. R. Jakobsen and K. H. Karlsen. A ”maximum principle for semicontinuous functions” applicable to integro-partial differential equations. *NoDEA Nonlinear Differential Equations Appl.*, 13:137-165, 2006.
- [18] E. R. Jakobsen, K. H. Karlsen, and C. L. Chioma. Error estimates for approximate solutions to Bellman equations associated with controlled jump-diffusions. Submitted to *Numer. Math.*, 2005.
- [19] N. V. Krylov. On the rate of convergence of finite-difference approximations for Bellman’s equations. *Algebra i Analiz*, 9(3):245–256, 1997.
- [20] N. V. Krylov. On the rate of convergence of finite-difference approximations for Bellman’s equations with variable coefficients. *Probab. Theory Related Fields*, 117(1):1–16, 2000.
- [21] N. V. Krylov. The rate of convergence of finite-difference approximations for Bellman equations with Lipschitz coefficients. *Appl. Math. Optim.*, 52(3):365–399, 2005.
- [22] R. Mikulyavichyus and G. Pragarauskas. Nonlinear potentials of the Cauchy-Dirichlet problem for the Bellman integro-differential equation. *Liet. Mat. Rink.*, 36(2):178–218, 1996.
- [23] H. Pham. Optimal stopping of controlled jump diffusion processes: a viscosity solution approach. *J. Math. Systems Estim. Control*, 8(1):27 pp. (electronic), 1998.

(Imran H. Biswas)

CENTRE OF MATHEMATICS FOR APPLICATIONS, UNIVERSITY OF OSLO, P.O. BOX 1053, BLINDERN, NO-0316 OSLO, NORWAY

*E-mail address:* [i.h.biswas@cma.uio.no](mailto:i.h.biswas@cma.uio.no)

(Espen R. Jakobsen)

NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, N-7491, TRONDHEIM, NORWAY

*E-mail address:* [erj@math.ntnu.no](mailto:erj@math.ntnu.no)

*URL:* [www.math.ntnu.no/~erj/](http://www.math.ntnu.no/~erj/)

(Kenneth Hvistendahl Karlsen)

CENTRE OF MATHEMATICS FOR APPLICATIONS, UNIVERSITY OF OSLO, P.O. BOX 1053, BLINDERN, NO-0316 OSLO, NORWAY

*E-mail address:* [kennethk@math.uio.no](mailto:kennethk@math.uio.no)

*URL:* [folk.uio.no/kennethk/](http://folk.uio.no/kennethk/)