

A uniformly converging scheme for fractal conservation laws

Jérôme Droniou and Espen R. Jakobsen

Abstract The fractal conservation law $\partial_t u + \partial_x(f(u)) + (-\Delta)^{\alpha/2} u = 0$ changes characteristics as $\alpha \rightarrow 2$ from non-local and weakly diffusive to local and strongly diffusive. In this paper we present a corrected finite difference quadrature method for $(-\Delta)^{\alpha/2}$ with $\alpha \in [0, 2]$, combined with usual finite volume methods for the hyperbolic term, that automatically adjusts to this change and is uniformly convergent with respect to $\alpha \in [\eta, 2]$ for any $\eta > 0$. We provide numerical results which illustrate this asymptotic-preserving property as well as the non-uniformity of previous finite difference or finite volume type of methods.

1 Introduction

We consider the following fractional conservation law

$$\begin{aligned} \partial_t u_\alpha + \partial_x(f(u_\alpha)) + \mathcal{L}_\alpha[u_\alpha] &= 0, \quad t > 0, x \in \mathbb{R}, \\ u_\alpha(0, x) &= u_{\text{ini}}(x), \quad x \in \mathbb{R}, \end{aligned} \quad (1)$$

where $\alpha \in [0, 2]$, $\mathcal{L}_\alpha = (-\Delta)^{\alpha/2}$,

$$u_{\text{ini}} \in L^\infty(\mathbb{R}) \cap BV(\mathbb{R}) \quad \text{and} \quad f: \mathbb{R} \rightarrow \mathbb{R} \text{ is locally Lipschitz-continuous.} \quad (2)$$

Such models appear for example in mathematical finance, gas detonation or semiconductor growth [23, 26, 11, 1]. The fractional Laplacian $\mathcal{L}_\alpha = (-\Delta)^{\alpha/2}$ can be

Jérôme Droniou

School of Mathematical Sciences, Monash University, Victoria 3800, Australia, e-mail: jerome.droniou@monash.edu

Espen R. Jakobsen

Department of Mathematical Sciences, Norwegian University of Science and Technology, 7491 Trondheim, Norway, e-mail: erj@math.ntnu.no

defined e.g. as a Fourier multiplier, but for our purpose the following equivalent definition, valid for any $\varphi \in C_c^\infty(\mathbb{R})$ (set of smooth compactly supported functions), is more useful:

$$\begin{cases} \mathcal{L}_0[\varphi](x) = \varphi(x), & \alpha = 0, \\ \mathcal{L}_\alpha[\varphi](x) = -c_\alpha \int_{\mathbb{R}} \frac{\varphi(x+z) - \varphi(x) - \varphi'(x)z \mathbf{1}_{[-1,1]}(z)}{|z|^{1+\alpha}} dz, & \alpha \in (0, 2), \\ \mathcal{L}_2[\varphi](x) = -\Delta \varphi(x), & \alpha = 2, \end{cases} \quad (3)$$

where $\mathbf{1}_{[-1,1]}$ is the characteristic function of $[-1, 1]$, $c_\alpha = (2\pi)^\alpha \frac{\alpha \Gamma(\frac{1+\alpha}{2})}{2\pi^{\frac{1}{2}+\alpha} \Gamma(1-\frac{\alpha}{2})}$ and Γ is the Euler function [15].

As $\alpha \rightarrow 2$, the operator \mathcal{L}_α changes nature and properties. For $\alpha \in (0, 2)$, \mathcal{L}_α is a *non-local* pseudo-differential operator of order < 2 , and it has relatively weak diffusive properties since the decay at infinity of the fundamental solution of $\partial_t u + \mathcal{L}_\alpha[u] = 0$ is polynomial. At $\alpha = 2$, $\mathcal{L}_\alpha = -\Delta$ is a *local* operator with strong diffusive properties and a fundamental solution with super-exponential decay. When α vary over $[0, 2]$, the qualitative behaviour of the solution u_α of (1) also changes. In the case that $\alpha = 2$, it is well-known that u_α becomes instantly smooth for $t > 0$ even when the initial data is discontinuous. On the contrary, for $\alpha = 0$, the solution may develop shocks and uniqueness of the solution requires additional entropy conditions and the corresponding notion of entropy solution [22]. The study of the fractional case $\alpha \in (0, 2)$ dates back to [6], with some restrictions on α and f . The first complete study in the case $\alpha > 1$ for any locally Lipschitz f and bounded initial data u_{ini} can be found in [14]. Here it is proved that the solution becomes instantly smooth even if u_{ini} is only bounded (see also [15]). If $\alpha < 1$, then the solution can develop shocks [4] and the weak solution need not be unique [3]. The notion of entropy solution of [2] is therefore required to obtain a well-posed formulation.

There exists a vast literature on the numerical approximation of scalar conservation laws (i.e. (1) without \mathcal{L}_α), see e.g. [17, 18, 19] and references therein. The study of numerical methods for fractal conservation laws is much more recent with a corresponding less extensive literature. Probabilistic methods have been studied in [21, 24], but must be applied to the equation satisfied by $\partial_x u_\alpha$ in order to avoid noisy results, and recovering from this a numerical approximation of u_α may be challenging in dimension greater than 1. Deterministic methods for (1) like finite difference, volume, and element methods (discontinuous Galerkin) are given in [13, 8, 10], while a high order spectral vanishing viscosity method is introduced in [9]. The latter method and its analysis is very different from the former three methods, with convergence and (non-optimal) error estimates that are independent of $\alpha \in (0, 2)$. As opposed to the spectral method, the other methods are monotone or have low order monotone variants.

Surprisingly, for all the non-spectral monotone methods the convergence deteriorates as $\alpha \rightarrow 2$, and the schemes themselves are not even defined in the limit $\alpha = 2$. The purpose of this paper is to present an *asymptotic-preserving monotone* scheme for (1) defined for any $\alpha \in [0, 2]$, i.e. a scheme that provides a monotone

approximation of u_α which is uniform with respect to $\alpha \in [0, 2]$. In particular, our scheme naturally adapts to the change of behaviour of \mathcal{L}_α as $\alpha \rightarrow 2$ and $\alpha \rightarrow 0$ and its convergence properties do not deteriorate in these extreme cases. The idea behind our scheme is to add a correction term in the form of a suitably chosen vanishing local viscosity term. Similar ideas have been used for other equations before, see e.g. [12] for linear equations and [20] for fully nonlinear equations. A stochastic interpretation can be found in [5].

This paper is organised as follows. The numerical method is presented in Section 2, and its asymptotic-preserving characteristics are discussed. Due to lack of space and the technical nature of the proofs, we skip them and refer instead to [16]. In Sections 3 and 4, we define precisely what asymptotic preserving means and the we give a couple numerical simulations to illustrate this property of the method.

2 The scheme

The new scheme is based on monotone conservative finite difference approximations of the local terms combined with quadrature, truncation of $\frac{1}{|z|^{1+\alpha}}$ near the singularity, and a second order correction term (vanishing viscosity) for the non-local term. Except for the correction term, the scheme is similar to the schemes of [13, 8] and of [10] with P_0 -elements. It is monotone, conservative, and converges in L^1_{loc} uniformly in $\alpha \in [\eta, 2]$ for all $\eta > 0$.

For given space and time steps $\delta x, \delta t > 0$, we introduce the grid $t_n := n\delta t$ and $x_i := i\delta x + \frac{\delta x}{2}$ for $n \in \mathbb{N}_0$ and $i \in \mathbb{Z}$. We identify sequences $(\varphi_i)_{i \in \mathbb{Z}}$ of numbers with piecewise constant functions $\varphi_{\delta x} : \mathbb{R} \rightarrow \mathbb{R}$ equal to φ_i on $[i\delta x, (i+1)\delta x)$ for all $i \in \mathbb{Z}$. Similarly, $(\varphi_i^n)_{n \geq 0, i \in \mathbb{Z}}$ is identified with $\varphi_{\delta x, \delta t} : [0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$ equal to φ_i^n on $[n\delta t, (n+1)\delta t) \times [i\delta x, (i+1)\delta x)$ for all $n \geq 0$ and $i \in \mathbb{Z}$. The discretisation of (1) can then be written as: find $u_{\alpha, \delta x, \delta t} = (u_i^n)_{n \geq 0, i \in \mathbb{Z}}$ such that

$$u_i^0 = \frac{1}{\delta x} \int_{[i\delta x, (i+1)\delta x)} u_0(x) dx \quad \text{for all } i \in \mathbb{Z}, \quad (4)$$

$$\frac{u_i^{n+1} - u_i^n}{\delta t} + \mathcal{F}_{\delta x}(u^n)_i + \mathcal{L}_{\alpha, \delta x}[u^{n+1}]_i = 0 \quad \text{for all } n \geq 0 \text{ and all } i \in \mathbb{Z}, \quad (5)$$

where $\mathcal{F}_{\delta x}$ is any monotone consistent and conservative discretization of $\partial_x(f(u))$ (see e.g. [17, 18, 19]), and $\mathcal{L}_{\alpha, \delta x}$ is a monotone discretisation of \mathcal{L}_α to be defined. Note that the scheme has explicit convection and implicit diffusion terms.

The first and simplest idea to obtain a monotone discretization of \mathcal{L}_α for $\alpha \in (0, 2)$ is to discretize the integral in (3) using a simple (weighted) midpoint type quadrature rule, see e.g. [13, 10, 8]. For $\varphi \in C_c^\infty(\mathbb{R})$ and letting $\varphi_l = \varphi(x_l)$ if $l \in \mathbb{Z}$, this leads to

$$\mathcal{L}_\alpha[\varphi](x_i) \approx \tilde{\mathcal{L}}_{\alpha, \delta x}[\varphi]_i := - \sum_{j \in \mathbb{Z} \setminus \{0\}} \left(\varphi_{i+j} - \varphi_i \right) \int_{(j\delta x - \frac{\delta x}{2}, j\delta x + \frac{\delta x}{2})} \frac{c_\alpha}{|z|^{1+\alpha}} dz. \quad (6)$$

However, as $\alpha \rightarrow 2$ we have $c_\alpha \rightarrow 0$ and therefore $\tilde{\mathcal{L}}_{\alpha, \delta x} \rightarrow 0$ for fixed δx . In the limit $\alpha \rightarrow 2$ the scheme then converges to

$$\frac{u_i^{n+1} - u_i^n}{\delta t} + \mathcal{F}_{\delta x}(u^n)_i = 0 \quad \text{for all } n \geq 0 \text{ and all } i \in \mathbb{Z},$$

which is a discretisation of $\partial_t u + \partial_x(f(u)) = 0$ and not $\partial_t u + \partial_x(f(u)) - \Delta u = 0$. Hence the limits $\alpha \rightarrow 2$ and $\delta x \rightarrow 0$ do not commute and the scheme is not asymptotic-preserving.

Note that $\tilde{\mathcal{L}}_{\alpha, \delta x}$ vanishes in the limit because the measure $\frac{c_\alpha dz}{|z|^{1+\alpha}}$ concentrates around 0 as $\alpha \rightarrow 2$, while in the above midpoint rule the integral in (3) over $(-\frac{\delta x}{2}, \frac{\delta x}{2})$ will always be zero by symmetry. We therefore need to replace the midpoint rule on this interval by a more accurate rule based on the second order interpolation polynomial P_i of φ around the node x_i . We find that this polynomial satisfies $P_i(x_i + z) - P_i(x_i) - P_i'(x_i)z = \frac{1}{2\delta x^2} (z^2\varphi_{i-1} - 2z^2\varphi_i + z^2\varphi_{i+1})$ and the new discretization therefore becomes

$$\begin{aligned} \mathcal{L}_{\alpha, \delta x}[\varphi]_i &:= -c_\alpha \int_{-\frac{\delta x}{2}}^{\frac{\delta x}{2}} \frac{P(x_i + z) - P(x_i) - P'(x_i)z}{|z|^{1+\alpha}} dz + \tilde{\mathcal{L}}_{\alpha, \delta x}[\varphi]_i \\ &= \frac{\varphi_{i+1} - 2\varphi_i + \varphi_{i-1}}{\delta x^2} \int_{(-\frac{\delta x}{2}, \frac{\delta x}{2})} \frac{c_\alpha |z|^{1-\alpha}}{2} dz + \tilde{\mathcal{L}}_{\alpha, \delta x}[\varphi]_i. \end{aligned}$$

We can check that the new approximation has the following truncation error [16]:

$$\begin{aligned} &|\mathcal{L}_\alpha[\varphi](x_i) - \mathcal{L}_{\alpha, \delta x}[\varphi]_i| \\ &\leq C \left(\|\varphi^{(4)}\|_{L^\infty} \delta x^{4-\alpha} + \|\varphi''\|_{L^\infty} c_\alpha \left(\frac{1}{\alpha} + \frac{1}{|1-\alpha|} \right) \delta x^{\min(1, 2-\alpha)} + \|\varphi'\|_{L^\infty} \delta x \right), \end{aligned}$$

which is $O(\delta x) + o_\alpha(1)$ as $\alpha \rightarrow 2$ and therefore does not deteriorate in this limit. Note that if $\alpha = 1$, then $\frac{1}{|1-\alpha|} \delta x^{\min(1, 2-\alpha)}$ must be replaced with $\delta x |\ln(\delta x)|$.

In order to obtain an approximation which uses only a finite number of discrete values, we also truncate the sum in (6) as in [13] at some index $J_{\delta x} > 0$ (which may depend upon α) where $J_{\delta x} \delta x \rightarrow \infty$ as $\delta x \rightarrow 0$. The final approximate operator $\mathcal{L}_{\alpha, \delta x}$ is therefore

$$\begin{aligned} \mathcal{L}_{\alpha, \delta x}[\varphi]_i &= - \sum_{0 < |j| \leq J_{\delta x}} W_{\alpha, \delta x}^j (\varphi_{i+j} - \varphi_i) - W_{\alpha, \delta x}^{J_{\delta x}+1} (\varphi_{i-J_{\delta x}-1} - \varphi_i) \\ &\quad - W_{\alpha, \delta x}^{J_{\delta x}+1} (\varphi_{i+J_{\delta x}+1} - \varphi_i) - W_{\alpha, \delta x}^0 \frac{\varphi_{i+1} - 2\varphi_i + \varphi_{i-1}}{\delta x^2}, \quad (7) \end{aligned}$$

with weights

$$\begin{aligned}
W_{\alpha, \delta x}^0 &= \int_{(-\frac{\delta x}{2}, \frac{\delta x}{2})} \frac{c_\alpha |z|^{1-\alpha}}{2} dz, \\
W_{\alpha, \delta x}^j &= \int_{(j\delta x - \frac{\delta x}{2}, j\delta x + \frac{\delta x}{2})} \frac{c_\alpha}{|z|^{1+\alpha}} dz \quad \text{for } 0 < |j| \leq J_{\delta x}, \\
W_{\alpha, \delta x}^{J_{\delta x}+1} &= \int_{z > J_{\delta x} \delta x + \frac{\delta x}{2}} \frac{c_\alpha}{|z|^{1+\alpha}} dz = \int_{z < -J_{\delta x} \delta x - \frac{\delta x}{2}} \frac{c_\alpha}{|z|^{1+\alpha}} dz.
\end{aligned} \tag{8}$$

The last term in (7) contains the classical discretization of $\varphi''(x_i)$ and is the new correction term compared with the discretisations of [13, 10, 8]. Discretisation (7)–(8) fits in the generic framework of [13] from which we can conclude:

Theorem 1 ([16]). *Under a standard CFL condition for the convection term,*

1. *There is a unique solution $u_{\alpha, \delta x, \delta t}$ of the scheme defined by (4), (5), (7) and (8), satisfying $\|u_{\alpha, \delta x, \delta t}\|_{L^\infty} \leq \|u_{ini}\|_{L^\infty}$ and $|u_{\alpha, \delta x, \delta t}(t, \cdot)|_{BV} \leq |u_{ini}|_{BV}$ for all $t > 0$.*
2. *For fixed α , $u_{\alpha, \delta x, \delta t}$ converges in $L_{loc}^1([0, \infty) \times \mathbb{R})$ as $(\delta x, \delta t) \rightarrow 0$ to the unique entropy solution u_α of (1).*

Remark 1. We set $\mathcal{L}_{2, \delta x}[\varphi]_i = -(\varphi_{i+1} - 2\varphi_i + \varphi_{i-1})/\delta x^2$ and $\mathcal{L}_{0, \delta x}[\varphi]_i = \varphi_i$. This consists in fixing δx and sending $\alpha \rightarrow 2$ or $\alpha \rightarrow 0$ in (7). Taking the limits in the scheme (5), we obtain the classical implicit scheme for the (1) with $\alpha = 2$ or $\alpha = 0$.

3 The asymptotic-preserving property

The scheme is asymptotic-preserving if its solution $u_{\alpha, \delta x, \delta t}$ satisfies the following uniform approximation result away from $\alpha = 0$ (see [16] for the case $\alpha = 0$):

$$\forall \eta > 0, \quad \sup_{\alpha \in [\eta, 2]} d_{L_{loc}^1([0, \infty) \times \mathbb{R})}(u_{\alpha, \delta x, \delta t}, u_\alpha) \rightarrow 0 \text{ as } (\delta x, \delta t) \rightarrow 0 \tag{9}$$

where $d_{L_{loc}^1([0, \infty) \times \mathbb{R})}(u, v) = \sum_{n=1}^{\infty} 2^{-n} \min(1, \|u - v\|_{L^1([0, n) \times (-n, n])})$ is the usual distance defining the topology of $L_{loc}^1([0, \infty) \times \mathbb{R})$. Here and elsewhere, the convergence $(\delta x, \delta t) \rightarrow 0$ is always taken under a standard CFL condition depending on the definition of the convective flux \mathcal{F} in (5) (see e.g. [13, 10, 8]). This formulation of the asymptotic-preserving property is very general and does not require an explicit error estimate independent on α . Such an estimate seems particularly challenging to obtain in the absence of regularity of the solution as $t \rightarrow 0$.

Theorem 2 ([16]). *Under a standard CFL for the convection part, the numerical scheme defined by (4) (5), (7) and (8) is asymptotic-preserving.*

Next we want to illustrate this property numerically. As it is formulated now, this would require us to have access to the exact solution u_α , which is not the case. We overcome this difficulty by using instead the following equivalent reformulation of (9) (see [16]), which can be checked by computing approximate solutions only:

$\forall \alpha_0 \in (0, 2]$, for any sequence $(\delta x_k, \delta t_k)_{k \in \mathbb{N}}$ converging to 0:

$$\sup_{k \geq 1} d_{L^1_{\text{loc}}([0, \infty) \times \mathbb{R})}(\mathbf{u}_{\alpha, \delta x_k, \delta t_k}, \mathbf{u}_{\alpha_0, \delta x_k, \delta t_k}) \rightarrow 0 \text{ as } \alpha \rightarrow \alpha_0. \quad (10)$$

Remark 2. The matrix of $\mathcal{L}_{\alpha, \delta x}$ defined by (7) is a semi-definite Toeplitz matrix as in [13, 10, 8]. Implementation of the scheme thus takes advantage of super-fast multiplication and inversion algorithms for these matrices [7, 25]. Computing several approximate solutions, as required in (10), is therefore not very expensive.

4 Numerical results

In all these tests, we take the Burgers flux $f(u) = \frac{u^2}{2}$ and $\mathcal{F}_{\delta x}$ given by a MUSCL method. The final time is $T = 1$ and the spatial computational domain is $[-1, 1]$. We use the same truncation parameters (in particular $J_{\delta x}$) as in [13, Section 4.1.2].

For each test, we choose the discretisation steps $(\delta x_k, \delta t_k) = (\frac{1}{2^k \times 50}, \frac{1}{2^k \times 100})$ for $k = 1, \dots, 4$, which all satisfy the CFL for (5). We also select four values $(\alpha_m)_{m=1, \dots, 4} = (1.8, 1.9, 1.99, 1.999)$ which are near $\alpha_0 = 2$, the difficult case in assessing the uniformity of the convergence in (10) and the reason why we introduced the correction term in (7). We then indicate, for $m = 1, \dots, 4$, the value of

$$E_m = \max_{k=1, \dots, 4} \sup_{t \in [0, 1]} \|u_{\alpha_m, \delta x_k, \delta t_k}(\cdot, t) - u_{\alpha_0, \delta x_k, \delta t_k}(\cdot, t)\|_{L^1([-1, 1])},$$

that is the maximum over $k = 1, \dots, 4$ of the $L^\infty(L^1)$ norm of $u_{\alpha_m, \delta x_k, \delta t_k} - u_{\alpha_0, \delta x_k, \delta t_k}$ on the computational domain. This is a stronger norm than the $L^1(L^1)$ norm used in (10). Hence, E_m approaching 0 as m increases is an even better indication that the scheme is asymptotic-preserving.

Test 1 (rarefaction): we select a Riemann initial condition, $u_{\text{ini}} = -1$ if $x < 0$ and $u_{\text{ini}} = 1$ if $x > 0$. In this case both convection and diffusion work to smooth out the initial data. Table 1 shows the values of $(E_m)_{m=1, \dots, 4}$ for both the uncorrected scheme from [13] based on (6) and our corrected scheme based on (7).

Table 1 Comparison between the uncorrected scheme of [13] and our corrected scheme, $u_{\text{ini}} = -1$ on $(-\infty, 0)$, $u_{\text{ini}} = 1$ on $(0, \infty)$.

	E_1	E_2	E_3	E_4
Uncorrected scheme	1.8E-1	3E-1	8.8E-1	9.1E-1
Corrected scheme	5.1E-2	2.2E-2	1.7E-4	1.7E-5

Test 2 (smooth shock): the initial condition is $u_{\text{ini}}(x) = 1$ if $x < 0$ and $u_{\text{ini}}(x) = -1$ if $x > 0$. Here the hyperbolic and non-local terms in (1) compete to maintain or diffuse the initial shock. Since α_m is near 2 however, any solution is instantly smooth, but has much larger gradients near $x = 0$ than the solution in Test 1 (Table 2).

Table 2 Comparison between the uncorrected scheme of [13] and our corrected scheme, $u_{\text{ini}} = 1$ on $(-\infty, 0)$, $u_{\text{ini}} = -1$ on $(0, \infty)$.

	E_1	E_2	E_3	E_4
Uncorrected scheme	2.1E-1	3.9E-1	1.3	1.3
Corrected scheme	5.3E-2	2.3E-2	3.2E-4	4.2E-5

Both tests confirm that the scheme defined by (4), (5), (7) and (8) is asymptotic-preserving. They also confirm that, without the order 2 correction in (7), the scheme deteriorates as $\alpha \rightarrow 2$ and does not provide a correct numerical solution at any reasonable resolution. This is also illustrated in Figure 1, where we plot the solutions of both schemes for $\alpha = 1.99$ for the initial condition of Test 2 and $(\delta x, \delta t) = (\frac{1}{2^4 \times 50}, \frac{1}{2^4 \times 100})$. Even at this very high resolution, the uncorrected scheme provides an incorrect approximate solution which, as expected, is closer to the solution of $\partial_t u + \partial_x(f(u)) = 0$ than to the solution of (1).




Fig. 1 Approximate solutions provided at $T = 1$ by the corrected (continuous) and uncorrected (dashed) schemes for (1) with $\alpha = 1.99$. The dotted line is both the initial condition and the solution to $\partial_t u + \partial_x(f(u)) = 0$.

5 Conclusion

We have presented a monotone numerical method for fractional conservation laws which is asymptotic-preserving with respect to the fractional power of the Laplacian. The scheme automatically adjusts to the change of nature of the equation as the power of the Laplacian goes to 1 (i.e. $\alpha \rightarrow 2$ in (1)) and therefore provides accurate approximate solutions for any power of the fractional Laplacian. We have given numerical results to illustrate the asymptotic-preserving property of our method, as well as the necessity of modifying previously studied monotone methods to obtain this property.

The complete theoretical study of such monotone asymptotic-preserving schemes will be presented in the forthcoming paper [16]. Here a general class of fractional degenerate parabolic equations are considered that include (1) as a special case.

References

1. Alfaro, M., Droniou, J.: General fractal conservation laws arising from a model of detonations in gases. *Appl. Math. Res. Express* **2012**, 127–151 (2012)
2. Alibaud, N.: Entropy formulation for fractal conservation laws. *J. Evol. Equ.* **7**(1), 145–175 (2007)
3. Alibaud, N., Andreianov, B.: Non-uniqueness of weak solutions for the fractal Burgers equation. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **27**(4), 997–1016 (2010)
4. Alibaud, N., Droniou, J., Vovelle, J.: Occurrence and non-appearance of shocks in fractal burgers equations. *J. Hyperbolic Differ. Equ.* **4**(3), 479–499 (2007)
5. Asmussen, S., Rosiński, J.: Approximations of small jumps of Lévy processes with a view towards simulation. *J. Appl. Probab.* **38**(2), 482–493 (2001)
6. Biler, P., Karch, G., Woyczynski, W.: Fractal burgers equations. *J. Diff. Eq.* **148**, 9–46 (1998)
7. Chan, R., Ng, M.: Conjugate gradient methods for toeplitz systems. *SIAM Review* **38**(3), 427–482 (1996)
8. Cifani, S., Jakobsen, E.R.: On numerical methods and error estimates for degenerate fractional convection-diffusion equations. To appear in *Numer. Math.* DOI 10.1007/s00211-013-0590-0
9. Cifani, S., Jakobsen, E.R.: On the spectral vanishing viscosity method for periodic fractional conservation laws. *Math. Comp.* **82**(283), 1489–1514 (2013)
10. Cifani, S., Jakobsen, E.R., Karlsen, K.H.: The discontinuous galerkin method for fractal conservation laws. *IMA J. Numer. Anal.* **31**(3), 1090–1122 (2011)
11. Clavin, P.: *Instabilities and nonlinear patterns of overdriven detonations in gases.* Kluwer (2002)
12. Cont, R., Tankov, P.: *Financial modelling with jump processes.* Chapman & Hall/CRC Financial Mathematics Series. Chapman & Hall/CRC, Boca Raton (FL) (2004)
13. Droniou, J.: A numerical method for fractal conservation laws. *Math. Comp.* **79**(269), 95–124 (2010)
14. Droniou, J., Gallouët, T., Vovelle, J.: Global solution and smoothing effect for a non-local regularization of an hyperbolic equation. *J. Evol. Equ.* **3**(3), 499–521 (2003)
15. Droniou, J., Imbert, C.: Fractal first order partial differential equations. *Arch. Ration. Mech. Anal.* **182**(2), 299–331 (2006)
16. Droniou, J., Jakobsen, E.R.: An asymptotic-preserving scheme for fractal conservation laws and fractional degenerate parabolic equations. In preparation
17. Eymard, R., Gallouët, T., Herbin, R.: Finite volume methods. In: P.G. Ciarlet, J.L. Lions (eds.) *Techniques of Scientific Computing, Part III, Handbook of Numerical Analysis, VII*, pp. 713–1020. North-Holland, Amsterdam (2000)
18. Godlewski, E., Raviart, P.A.: Numerical approximation of hyperbolic systems of conservation laws, *Applied Mathematical Sciences*, vol. 118. Springer, New-York (1996)
19. Holden, H., Holden, R.N.: *Front tracking for Hyperbolic Conservation Laws.* Springer (2002)
20. Jakobsen, E.R., Karlsen, K.H., La Chioma, C.: Error estimates for approximate solutions to Bellman equations associated with controlled jump-diffusions. *Numer. Math.* **110**(2), 221–255 (2008)
21. Jourdain, B., Méléard, S., Woyczynski, W.: Probabilistic approximation and inviscid limits for one-dimensional fractional conservation laws. *Bernoulli* **11**(4), 689–714 (2005)
22. Kruzhkov, S.N.: First order quasilinear equations with several independent variables. *Math. Sb. (N.S.)* **81**(123), 228–255 (1970)
23. Soner, H.: Optimal control with state-space constraint ii. *SIAM J. Control Optim.* **24**(6) (1986)
24. Stanescu, D., Kim, D., Woyczynski, W.: Numerical study of interacting particles approximation for integro-differential equations. *Journal of Computational Physics* **206**, 706–726 (2005)
25. Van Loan, C.: *Computational frameworks for the fast Fourier transform, Frontiers in Applied Mathematics*, vol. 10. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA (1992)
26. Woyczynski, W.: *Lévy processes in the physical sciences.* Birkhäuser, Boston (2001)